

PROCEEDINGS  
OF THE  
ZOOLOGICAL SOCIETY  
OF LONDON.



---

PART XXIII.

1855.

---

PRINTED FOR THE SOCIETY ;  
SOLD AT THEIR HOUSE IN HANOVER-SQUARE,  
AND BY MESSRS. LONGMAN, BROWN, GREEN AND LONGMANS,  
PATERNOSTER-ROW.



FIG. X. *Conularia*. 2. *almae* Macbrayne. (Berk. var. *aterrima* + *darkalis* C. Gmelin.) *Verreauxi* (F. E. Muller) *var. *maculata** (A. Nodder) (D. Draparnaud). *Reticulata* (Lamarck).

*gine dextro subrepando, columellari paulo breviore, dilatato, patente.*

Long. 15–17, diam.  $7\frac{3}{4}$ – $8\frac{1}{4}$  mill.

Hab. — ?

47. ENNEA REEVEANA, Pfr. *E. testa breviter rimata, cylindracea, tenuiscula, oblique distincte plicato-striata, nitida, virenti-hyalina; spira subcylindrica, apice conoideo-rotundata; sutura impressa, submarginata, ad anfr. ultimum conferte denticulata; anfr. 7, primis 2 lœvigatis, sequentibus convexiusculis, ultimo  $\frac{3}{7}$  longitudinis formante, antice subascendente, basi rotundato; columella oblique recedente, compressa; apertura subverticali, acuminato-ovali; perist. albo, reflexiusculo, margine dextro subrepando, columellari breviore, sursum dilatato, fere adnato.*

Long. 15, diam. 6 mill.

Hab. — ?

5. DESCRIPTIONS OF A NEW GENUS AND TWENTY-THREE NEW SPECIES OF PNEUMONOPOMA, FROM THE COLLECTION OF H. CUMING, ESQ. BY DR. L. PFEIFFER.

(Mollusca, Pl. XXXII.)

1. TROCHATELLA EXCAVATA, Pfr. *Tr. testa depresso-globosa, spiraliter levissime lirata, opaca, carnea, convexa, acute mucronata; anfr. 6 vix convexiusculis, ultimo inflato, obsoletissime subangulato; columella lata, triangulari, excavata; apertura diagonali, ampla, subsemicirculari, in fundo fusco-carnea; perist. albo, late expanso, ad columellæ angulum subrotundatum angustato.*

Diam. maj.  $13\frac{1}{2}$ , min.  $11\frac{1}{2}$ , alt.  $9\frac{1}{2}$  mill.

Hab. — ?

2. HELICINA REPANDA, Pfr. *H. testa conoideo-semiglobosa, solida, levissime striatula, nitidula, lutescenti-albida, fascia 1 rubra supramediana ornata; spira convexa, subacuminata; anfr.  $4\frac{1}{2}$  vix convexiusculis, ultimo antice descendente; columella brevi, simplice, callum crassum, circumscriptum retrorsum emitente; apertura diagonali, subtriangulari; perist. recto, in-crassato, margine supero repando, basali versus columellam attenuato.*

Diam. maj. 8, min.  $6\frac{1}{2}$ , alt. 5 mill.

Hab. — ?

3. HELICINA ZEBRIOLATA, Pfr. *H. testa subconoideo-depressa, tenui, striatula, parum nitida, strigis flexuosis rubris et albis variegata, vel rubra, albo-punctata; spira parvula, parum elata, subacuta; anfr.  $4\frac{1}{2}$  planis, ultimo lato, acute carinato, basi inflato; columella brevi, simplice, callum emittente tenuem; aper-*

*tura perobliqua, semiovato-subtriangulari; perist. simplice, recto, margine basali in columellam subangulatim transeunte.*

Diam. maj.  $6\frac{3}{4}$ , min.  $5\frac{1}{2}$ , alt.  $3\frac{1}{2}$  mill.

Hab. Lord Howe's Island, Australian Seas (*Macgillivray*).

4. **HELICINA RUDIS**, Pfr. *H. testa conoideo-depressa, solidula, ruditer striatula et submalleata, filo-carinata, fulva; spira conoidea, acutiuscula; anfr.  $4\frac{1}{2}$  planis, ultimo descendente, basi parum convexo; columella brevi, simplice, callum tenuem, circumscripsum retrorsum emittente; aperturā diagonali, subtriangulari; perist. simplice, recto, acuto, margine basali cum columella angulum subrectum formante.*

Diam. maj.  $4\frac{1}{3}$ , min.  $3\frac{1}{3}$ , alt. 3 mill.

Hab. —?

5. **HELIX SHANGHAIENSIS**, Pfr. *H. testa depresso conoidea, tenui, carinata, lavigata, pallide cornea, rufo-strigata; spira breviter conoidea, obtusula; anfr. 5 convexiusculis, ultimo non descendente, ad suturam et carinam subacutam pallide articulato, basi convxiore; columella brevi, verticali, callum emittente tenuem; apertura fere diagonali, subtriangulari; perist. simplice, recto, acuto, margine basali leviter arcuato, cum columella angulum formante.*

Diam. maj.  $4\frac{1}{2}$ , min. 4, alt.  $2\frac{2}{3}$  mill.

Hab. Shanghai, China (*Mr. Fortune*).

6. **HELICINA SPHÆROIDEA**, Pfr. *H. testa subglobosa, solidula, sub lente minutissime striata, parum nitente, pallide virenti-striatula; spira rotundata, subacuminata; anfr.  $4\frac{1}{2}$  convexiusculis, ultimo magno, infra medium obsoletissime angulato; columella albo-callosa, recedente, basi indistincte dentata, callum emittente diffusum; apertura diagonali, late semiovali; perist. simplice, brevissime expanso, margine basali cum columella angulum subacutum formante.*

Diam. maj. 7, min. 6, alt.  $5\frac{1}{2}$  mill.

Hab. Loyalty Island, Australian Seas (*Macgillivray*).

7. **HELICINA MERDIGERA**, Sallé MSS. *H. testa globoso-turbinate, tenui, irregulariter et subdistanter striata, sub tegumento lutoso rubicunda vel hyalina; spira elata, conica, acutiuscula; anfr.  $5\frac{1}{2}$  convexiusculis, ultimo utrinque convxiore; columella brevi, basi extrorsum subdentata, callum emittente tenuissimum; apertura diagonali, subsemicirculari; perist. simplice, breviter expanso, margine basali sensim in denticulum columellæ abeunte.*

Diam. maj. 6, min. 5, alt.  $4\frac{1}{2}$  mill.

Hab. Vera Cruz, Mexico (*Sallé*).

8. **HELICINA MACGILLIVRAYI**, Pfr. *H. testa subconoideo-depressa, solidula, spiraliter obsolete striata, nitidula, carnea vel pallide lilacea, indistincte fasciata; spira breviter conoidea, subacuminata; anfr.  $4\frac{1}{2}$  planis, ultimo peripheria obsoletissime angulato;*

*apertura perobliqua, subtriangulato-semiovali ; columella brevi, callum emittente crassiusculum, sulco brevi circumscriptum ; perist. simplice, breviter expanso, versus columellam attenuato.*

Diam. maj.  $6\frac{1}{2}$ , min.  $5\frac{1}{2}$ , alt. 4 mill.

Hab. Isle of Pines, Australian Seas (*Macgillivray*).

9. **HELICINA DENTICULATA**, Pfr. *H. testa globoso-conica, solidula, oblique striata, minutissime decussata, rugis antrorsum descendantibus distantioribus sculpta, parum nitida, alba, superne late flavo-fasciata ; spira convexo-conica, acuta ; sutura eleganter denticulata ; anfr.  $5\frac{1}{2}$  vix convexiusculis, ultimo peripheria subangulato, basi purum convexo ; columella breviter arcuata, antrorsum dentata, superne rima brevi notata ; retrorsum in callum tenuem abeunte ; apertura subtriangulari-semiovali ; perist. albo, angulatim patente, medio in rostrum triangulare producto.*

Diam. maj. a rostro 13, min. fere 11, alt. 9 mill.

Hab. Honduras.

10. **HELICINA TURBINELLA**, Pfr. *H. testa turbinata, solida, leviter striatula, vix nitidula, lutescente ; spira conoidea, acutiuscula ; anfr.  $4\frac{1}{2}$  convexiusculis, ultimo carinato, basi convexo ; columella brevi, subverticali, basi subtruncata, callum retrorsum emittente tenuissimum ; apertura diagonali, subtriangulari-semiovali ; perist. albido, tenui, breviter expanso, margine basali leviter arcuato.*

Diam. maj.  $4\frac{1}{3}$ , min. 4, alt. 3 mill.

Hab. Sidney.

11. **CYCLOSTOMA (CYCLOTUS) GLAUCOSTOMUM**, Pfr. *C. testa umbilicata, subturbinato-depressa, solidula, striata et plicis obliquis superne irregulariter sculpta, nitida, castanea ; spira brevissime conoidea, apice obtusulo, nudo ; anfr. 4 convexis, rapide accrescentibus, ultimo antice dilatato, brevi minute malato-granulata ; umbilico pervio,  $\frac{1}{4}$  diametri non æquante ; apertura vix obliqua, subrotundata, intus cœrulea, margaritacea ; perist. recto, marginibus superne subangulatim junctis, columellari subincrassato, perarcuato,*

Diam. maj. 18, min. 14, alt. 9 mill.

Hab. Venezuela.

12. **CYCLOSTOMA (CYCLOTUS) SORDIDUM**, Pfr. *C. testa umbilicata, conoideo-depressa, solidula, levissime et confertim spiraliter lirata, sub epidermide decidua, fulva sordide alba ; spira conoidea, mucronata ; anfr. 4 convexis, celeriter accrescentibus, ultimo terete, antice vix adnato ; apertura parum obliqua, subcirculari ; perist. simplice, continuo, vix expansiusculo.*

Diam. maj.  $7\frac{1}{2}$ , min.  $6\frac{1}{3}$ , alt.  $4\frac{1}{2}$  mill.

Hab. China.

13. **CYCLOSTOMA (CYCLOTUS) MACGILLIVRAYI**, Pfr. *C. testa umbilicata, turbinato-depressa, solida, rugoso-striata et liris*

*spiralibus in anfr. ultimo evanescentibus sculpta, opaca, carneo-albida; spira convexo-conoidea, submucronata; anfr. 4½ convexiusculis, sensim accrescentibus, ultimo subdepresso, basi circa umbilicum conicum,  $\frac{1}{3}$  diametri fere æquantem, abrupte angulato; apertura obliqua, superne et basi subangulata; perist. continuo, obtuso, recto, ad angulum circa umbilicalem subproducto.—Operc. concavo, calcareo, arctispirali.*

Diam. maj. 9, min.  $7\frac{1}{2}$ , alt. 5 mill.

Hab. Isle of Aneiteum, New Hebrides (*Macgillivray*).

14. **CYCLOSTOMA (AULOPOMA) GRANDE**, Pfr. *C. testa umbilicata, depressa, solida, confertim distincte striata, sulcis spiralibus nonnullis remotis sculpta, sub epidermide nigro-fusca, sæpe fasciatim detrita, obsolete marmorata; spira brevissima conoidea, vertice subtili, acuminato, nigricante; anfr. 5 convexis, ultimo terete, basi læviore, antice breviter soluto, vix descendente; apertura obliqua, subcirculari; perist. simplice, recto.—Operc. subsexspirato, anfractibus oblique valide plicatis.*

Diam. maj. 25, min.  $21\frac{1}{2}$ , alt. 13 mill.

Hab. Ceylon (*Mr. Thwaites*).

15. **CYCLOSTOMA (CYCLOPHORUS) FORBESIANUM**, Pfr. *C. testa umbilicata, subdepressa, solida, liris spiralibus inæqualibus subconfertis cincta, opaca, albida vel rubella; spira parum elevata, vertice submucronato; anfr. 5 convexis, ultimo descendente, terete; umbilico conico,  $\frac{1}{4}$  diametri superante; apertura parum obliqua, subcirculari, intus alba vel vinosa; perist. simplice, recto, superne subdilatato, breviter adnato.—Operc. pallido.*

Diam. maj. 23, min. 18, alt. 11 mill.

Hab. Lord Howe's Island; New Hebrides (*Macgillivray*).

16. **CYCLOSTOMA (CYCLOPHORUS) CONVEXIUSCULUM**, Pfr. *C. testa umbilicata, convexiuscula, tenuiuscula, epidermide sordide fuscescente, confertim membranaceo-plicata, ad suturam squamosa, induta; spira parum elevata, vertice subtili; anfr. 4½ modice convexis, ultimo subdepresso, basi rotundato; umbilico aperto,  $\frac{1}{3}$  diametri formante; apertura obliqua, subcirculari; perist. continuo, breviter adnato, simplice, recto.—Operc. corneo, indistincte arctespirato.*

Diam. maj. 6, min. 5, alt.  $2\frac{2}{3}$  mill.

Hab. Cape of Good Hope (*Macgillivray*).

17. **CYCLOSTOMA (LEPTOPOMA) BARBATUM**, Pfr. *C. testa umbilicata, depresso-turbinata, tenui, acute carinata, sub epidermide arcuato-striata, fulvo et castaneo variegata; spira turbinata, apice acuta; anfr. 5 convexiusculis, superne liris 3–4 prope carinam munitis, ultimo ad carinam pilis squamosis barbato, basi convexiore, liris 2 distantibus nonnullisque obsoletioribus subangulato; umbilico mediocri, conico; apertura fere diagonali, subcirculari; perist. duplice; interno continuo, expansiusculo, ad anfr. penul-*

*tum subsinuato, externo breviter patente et reflexiusculo, superne subdilatato.*

Diam. maj. 13, min. 11, alt.  $6\frac{1}{2}$  mill.

$\beta$ . *Major, aufr. 5, paulo depressior, umbilico paulo latiore.*

Diam. 18, alt.  $8\frac{1}{2}$  mill.

$\gamma$ . *Minor, anfr. 5, liris basalibus distinctioribus.*

Diam. 11, alt. 6 mill.

*Hab.* Borneo, Sarawak.

18. CYCLOSTOMA (CYCLOSTOMUS?) ANNATONENSE, Pfr. *C. testa perforata, globoso-conica, tenui, carinis pluribus periphericis acutis, lirisque compressis, confertis, vix elevatis munita, parum nitida, fusca; spira acute conica, sursum pallescente; anfr. 6 convexiusculis, ultimo ventroso; apertura vix obliqua, irregulariter ovali; perist. tenui, albido, marginibus approximatis, dextro brevissime expanso, columellari superne recedente, cum parte inferiore subangulatim juncto, angustissimo.*

Diam. maj. 8, min.  $6\frac{2}{3}$ , alt. 8 mill.

*Hab.* Isle of Aneiteum or Annaton; New Hebrides (*Macgillivray*).

19. CYCLOSTOMA (CYCLOSTOMUS?) SINUATUM, Pfr. *C. testa umbilicata, globoso-turbinata, solida, laevigata, vix nitidula, alba; spira turbinata, acutiuscula; anfr.  $5\frac{1}{2}$  convexis, prope suturam liris 5 filaribus cinctis, ultimo rotundato, circa umbilicum angustum, pervium liris paulo confertioribus sculpto; apertura vix obliqua, subcirculari; perist. breviter expanso, marginibus callo subexciso juncitis, dextro superne repando, fere sinuoso, columellari non latiore.*

Diam. maj. 22, min. 18, alt. 17 mill.

*Hab.* Madagascar?

20. DIPLOMATINA CHORDATA, Pfr. *D. testa sinistrorsa, profunde rimata, subfusiformi, tenui, laevigata, costulis subdistantibus chordæformibus munita, diaphana, cereo-albida; spira ovato-conica, acuta; anfr. 7 convexis, ultimo attenuato, antice ascendente; apertura vix obliqua, subcirculari; perist. duplice, interno continuo, breviter adnato, cæterum expansiusculo, externo breviter patente.*

Long. 4, diam. 2 mill.

$\beta$ . *Minor, paulo confertius chordata.*

*Hab.* New Zealand (*Strange*); var.  $\beta$ . Lord Howe's Island (*Macgillivray*).

ANAULUS, Pfr., nov. gen. *Operculum tenuissimum, corneum, subarctispirum, extus concaviusculum. Testa umbilicata, pupinæformis; apertura circularis; peristoma duplex: internum continuum, externum dilatum, ad insertionem marginis dextri canali aperto perforatum.*

21. ANAULUS BOMBYCINUS, Pfr. *A. testa impervie umbilicata, oblonga, tenuiuscula, confertim striata, sericea, rubello-cornea;*

*spira irregulari, inflata, in conum obtusulum desinente; anfr. 6 convexis, penultimo gibbo, latere aperturali subplanato, ultimo attenuato, ad suturam marginato, antice ascende; apertura circulari, basi axin subexcedente; perist. interno calloso, adnato, externo late expanso, reflexiusculo, marginibus superne callo junctis.*

—*Operc. luteo-corneo.*

Long. 14, diam.  $7\frac{1}{3}$  mill. Apert. intus fere 4 mill. diam.

$\beta.$  *Minor anfr. 5\frac{1}{2}, penultimo magis turgido.*

Long. 10, diam. vix 6 mill.

*Hab.* Borneo, Sarawak. Pl. XXXII. fig. 10.

22. HYDROCENA ACUTILIRATA, Pfr. *H. testa aperte perforata, turbinata, tenui, liris acutis, magis minusve confertis, nonnullis interdum carinæformibus, cincta, cornea; spira conica, acuta; anfr. 5\frac{1}{2} convexiusculis, ultimo basi inflato; apertura obliqua, angulato-subovali; perist. simplice, subrecto, marginibus vallo tenui junctis, sinistro subangulato.*

Long. 10, diam.  $6\frac{1}{2}$  mill.

*Hab.* Lord Howe's Island; New Hebrides (*Macgillivray*).

23. HYDROCENA (OMPHALOTROPIS) VESTITA, Pfr. *H. testa perforata, oblongo-conica, tenui, striata et spiraliter conferte lirata, epidermide fuscula vestita; spira pyramidata, acutiuscula; anfr. 6 planis, ultimo carina acuta infra medium, secunda circa perforationem, nonnullisque minoribus basalibus munita; apertura vix obliqua, subangulato-ovali; perist. simplice, recto, marginibus convergentibus.—Operc. fusco.*

Long. 5, diam. 3 mill. Ap. 2 mill. longa.

*Hab.* New Zealand.

## 6. DESCRIPTIONS OF NINE NEW SPECIES OF HELICEA, FROM MR. CUMING'S COLLECTION. BY DR. L. PFEIFFER.

(Mollusca, Pl. XXXII.)

1. SPIRAXIS CUMINGIANA, Pfr. *S. testa turrita, solida, sub-lævigata, nitidula, nigra; spira elongata, sursum in conum acutiusculum terminata; anfr. 8, supremis planis, capillaceo-striatis, reliquis prope suturam albo cinctam angulosis, ultimo  $\frac{2}{7}$  longitudinis subæquante, superne et infra medium angulato, basi attenuato; columella lamina alba, torta, basin fere attingente, munita; apertura vix obliqua, rhomboe-ovali, intus cærulescens; perist. simplice, intus nigro-labiato.*

Long. 48, diam. 16 mill.

*Hab.* Kanai, Sandwich Islands (*Dr. Newcomb*). Pl. XXXII. fig. 1.

2. BULIMUS TRAILLI, Pfr. *B. testa imperforata, oblongo-ovata, tenuiuscula, leviter striata, nitida, nigro-castanea; spira convexo-conica, apice obtusa; sutura levi, albo-marginata; anfr. 5 con-*