

PROCEEDINGS

OF THE

ZOOLOGICAL SOCIETY  
OF LONDON.



---

PART X.

1842.

---

PRINTED FOR THE SOCIETY,

BY R. AND J. E. TAYLOR, RED LION COURT, FLEET STREET.

*d.* Shell larger, with a pale band round the circumference, and a dark one beneath. Found under decayed leaves at Sinait, and in earth at Bolinao.

*e.* Shell very darkly coloured, size of *d.* On leaves of bushes at Calauang, and on leaves of trees at St. Christoval, in the province of Batangas.

*f.* Shell smoother and paler than the other varieties; apex blackish. Found on leaves of bushes at Daleguete, island of Zebu.

*g.* Shell larger than the other varieties, dark brown, with a white and brown articulated band close to the suture, and a nearly white circumferential band. Found on leaves of bushes at the island of Luban.

#### CYCLOSTOMA ALTUM.

*Cycl. testā acuminato-turritā, crassiusculā, fuscā, lāvi, tenuissimē striatā, apice obtusiusculo, anfractibus 7 ad 8 rotundatis; suturā distinctā, tenui; aperturā circulari, labio subincrassato, reflexo-patente, dupli, margine externo magis, interno minus expanso, intūs canālī parvā ad basin columellarem munito; umbilico mediocri extūs carinā obtusā marginato; operculo corneo, tenui, multi-spirali.* Long. 1.; lat. 0·3 poll.

*Hab.* supra truncos arborum in montibus insulae Negros, Philippinarum.

This species is remarkable for having a double lip, the inner or newer portion of which is not quite so much expanded as the outer. It may be regarded as the first link of affinity, connecting *Cyclostoma* with *Pupina* by the intervention of the next species, *C. Pupiniforme*.

#### CYCLOSTOMA PUPINIFORME.

*Cycl. testā subcylindrico-turritā, crassiusculā, obscurā, fuscā, lāvi, tenuissimē striatā, apice subabruptē acuminato, anfractibus senis, rotundatis, suturā validā; aperturā circulari, peritremate discontinuo; labio externo subincrassato, revoluto, flavido, ad basin columellarem canālī angustissimā spirali interrupto; deinde supernē latiori, demūn supra ultimum anfractum leviter expanso, canālī angustā tenuique inter anfractum ultimum et partem posticam labii posita; umbilico parvo, carinā obtusā marginato, hāce carinā externā partem canalis angustissimā basis columellaris efformante.* Long. 0·7; lat. 0·25.

*Hab.* supra truncos arborum apud S. Juan provinciæ Cagayan insulæ Luçon, Philippinarum.

This species, which is related to the last and to *Cycl. tortuosum* of Gray, approaches very nearly to the genus *Pupina*, appearing to differ only in having a dull unpolished external surface, while that of *Pupina* is extremely brilliant. It proves the genera *Cyclostoma* and *Pupina* to belong to the same family.

The next communication is from Dr. L. Pfeiffer, and contains the following descriptions of shells belonging to the genera *Helix* and *Bulimus*, also collected by H. Cuming, Esq. in the Philippine Islands.

**HELIX ZONIFERA**, Sow. *Hel. T. imperforatâ, subglobosâ, tenui, fulvâ, zoniis variis opacis lutescenti-albidis ornatâ, obsoletè angulatâ; anfractibus 4 convexis, supremis depressis; columellâ planâ, rectâ, elongatâ; aperturâ lunato-rotundatâ, intus nitidâ; peristomate simplici, expanso, albo, cum callo columellari angulatim juncto.*  
Diam. 1·60; alt. 1·10 poll.

*Hab.* ins. Leyte.

$\beta$ . *T. tenuissimâ, luteo-virente, basi unicolo, ad peripheriam fasciâ latâ nigricante et angustiore albida, supernè fasciis pluribus albidis interruptis ornatâ.*

*Hab.* ins. Leyte.

$\gamma$ . *T. rufo-nigricante, fasciis variis sordidè albidis.*

*Hab.* ins. Samar.

$\delta$ . *T. crassiusculâ, cinnamomeâ, fasciis lutescenti-albidis.*

*Hab.* ins. Samar.

Differt ab *H. pulcherrimâ*, cui valde affinis, spirâ depressâ, anfractu ultimo minus inflato, et columellâ neque excavatâ neque dentatâ.

**HELIX NORRISII**, Sow. *Hel. T. imperforatâ, globoso-depressâ, solidâ, glabra, fulvo-citrina, zonis opacis candidis et infra suturam maculis irregularibus albis notatâ; spirâ semiglobosa; anfractibus 4 $\frac{1}{2}$  convexis, ultimo ad columellam subexcavato; columella latâ, albo-callosâ, arcuatim prolongatâ; aperturâ ferè orbiculari, intus lactea; peristomate crasso, latissimè expanso.*

Diam. 1·80; alt. 1·15 poll.

*Hab.* insula Luçon, ad Montem Triga.

Species intermedia inter *H. pulcherrimam* et *zoniferam*, ab illâ spirâ depressâ et columellâ arcuatâ, ab alterâ columellâ excavatâ, nec angulatâ diversa.

**HELIX LUZONICA**, Sow. *Hel. T. imperforatâ, conico-globosâ, crassiusculâ, obliquè striatâ, rufâ, apice sanguineâ, epidermide libera ferè omnino tectâ, medio fasciâ latâ albida ornata; spirâ conoidea; anfractibus 5—5 $\frac{1}{2}$  convexiusculis; columella obliquâ, dilatata, tuberculosa; aperturâ lunato-orbiculari, intus lactea; peristomate incrassato, latè reflexo, albo vel purpureo latè limbato.*

Diam. 1·60; lat. 1·15 poll.

*Hab.* Provincia Cagayan insulæ Luçon.

Differt ab *H. pulcherrimâ* testâ crassâ, conoidea, anfractibus pluribus et sensim accentibus, columellâ vix excavatâ et labro incrassato.

**HELIX MINDANAENSIS**, Sow. *Hel. T. imperforatâ, globoso-conicâ, solidâ, obliquè rugoso-striatâ, apice obtuso, pallidâ, luteo-fuscâ, opacâ, maculis variis rufis pellucidis ornata; spirâ conoidea; anfractibus 4 $\frac{1}{2}$  convexiusculis, ultimo medio obtuse angulato, infra angulum fasciâ latâ, hydrophanâ, albida signata; columellâ livida, subrecte descendente, medio subintorta; aperturâ tetragono-ovatâ, intus plumbeâ; peristomate subincrassato, parum reflexo, livido-fusco, margine supero arcuatim dilatato.*

Diam. 2·20; alt. 1·65 poll.

*Hab.* insula Mindanao.

Forma et magnitudo sicut *H. Pan.* Differt anfractu ultimo angulato, colore et formâ columellæ et aperturæ. Ab *H. Harfordii* differt formâ conicâ, maculis non hydrophanis, sed in ipsâ testâ positis, et margine dextro aperturæ sinuato.

**HELIX CARBONARIA**, Sow. *Hel. T. imperforatâ, subtrocchiformi, lavi, purpureo-nigricante, epidermide fuscâ, hydrophanâ, obliquè strigata, apice rubrâ vel violacea; suturâ linearî; anfractibus 5 planiusculis, ultimo carinato, basi plano; columellâ subrectâ, dilatata, fuscidulâ; aperturâ subtetragonâ, intus livida; peristomate simplici, vix incrassato.*

Diam. 1·20; altit. 0·95 poll.

*Hab.* Insula Zebu. ‘Daleguete.’

Variat carinâ obtusiore, basi convexiore.

**HELIX (CAROCOLLA) PANAYENSIS**, Brod. *Hel. T. imperforatâ, depresso, orbiculari, carinatâ, crassisculâ, supernè griseo-fuscâ, minutissimè granulatâ, basi radiatim striatâ, nitidissimâ, olivaced; spirâ depresso-conoidea; anfractibus 6 planulatis, ultimo non deflexo; aperturâ angulato-lunari, intus albida; peristomate supernè simplici, basi incrassato, ad columellam expansiusculo.*

Diam. 1·60; altit. 0·85 poll.

*Hab.* Insula Panay. ‘Dingle.’

Var. (Cagayan ins. Luçon). *Spirâ elatiore, anfractibus ultimis tenuidis, supernè saturatè rufa, basi nigricans, deorsum pallescens.*

**HELIX MORICANDI**, Sow. *Hel. T. umbilicata, semiglobosa, basi planiuscula, nitida, albido-flavâ, fasciis rufis 2–3-cincta; anfractibus 5½ convexis, ultimo margine dextro subito deflexo; umbilico angusto, pervio; aperturâ transversè pyriformi, basi parallelâ; peristomate connexo, margine superiore expanso, basali latè reflexo, ad basin dente unico crasso instructo.*

Diam. 1·35; altit. 0·70 poll.

*Hab.* Insula Bohol. ‘Jacna.’

Differt ab *H. zonali*, Fér., cui persimilis, basi planâ, aperturæ parte supremâ deflexâ, indè horizontali, et marginibus peristomatis junctis.

**HELIX SAGITTIFERA**, Pfr. an Nanina? *Hel. T. subperforata, tenui, pellucidâ, obliquè striata et obsoletè rugosâ, fulvâ, maculis seriatis sagittiformibus et ad carinam obsoletam fasciâ unicâ rufâ ornata; suturâ impressâ, ad anfractum ultimum subcanaliculatâ; anfractibus 4½ planiusculis, ultimo inflato; aperturâ perobliquâ, lunato-ovatâ, intus lactea; peristomate simplici, ad columellam subincrassato, vix reflexo, margine superiore deflexo.*

Diam. 2; altit. 1·10 poll.

*Hab.* Sinait insulæ Luçon.

Affinis *H. Lamarckiana*, Lea. Differt formâ minus depressâ, anfractibus spiræ lente crescentibus, ultimo vix carinato, et colore.

Varietas: testâ distinctè carinatâ, supernè intensè rufâ, infra

carinam fasciâ nigricante dilutâ circumdatâ, basi olivaceo-fulgâ, maculis sagittiformibus obsoletis.—Bolinao insulæ Luçon.

**HELIX FULVIDA**, Pfr. an Nanina? *Hel. T. subperforatâ, subglobosâ, tenui, pellucidâ, pallidè fulvescente, supernè confertissimè et minutissimè granulosâ, basi glabrd, nitidâ; spirâ elatâ; anfractibus 5½, suprenis planis, 1½ ultimis rotundatis; aperturâ lunari; peristome simplici, margine columellari subincrassato, ad perforationem obsoletam reflexo.*

Diam. 1·25; altit. 0·85 poll.

Hab. Insula Mindanao.

**HELIX JANUS**, Chemn. xi. 3016. 17.—*Helicella, Fér. pr. 233.*—An Nanina? *Hel. T. sinistrorsâ, perforatâ, orbiculari, tenui, dia-phânâ, obliquè regulariter et confertim striatâ, supernè fuscâ, basi convexâ, rufâ, nitidiusculâ; spirâ latè depresso-conoideâ; anfractibus 7 planulatis, ultimo carinato; aperturâ lunari; peristome tenui, acuto, margine columellari reflexiusculo.*

Diam. 1·30; altit. 0·80 poll.

Hab. in monte Ophir, peninsulæ Malaccanæ.

*H. Mackensiana*, Saul in Revue Zool. 1841, p. 347, eadem esse videtur.

**HELIX PORPHYRIA**, Pfr. an Nanina? *Hel. T. perforatâ, depressâ, solidâ, obliquè rugoso-striatâ, rufâ, maculis et strigis creberrimis, flavidô-albidis sub prominulis marmoratâ, carinatâ, infra carinam rufo-fasciatâ, basi olivaceo-fulgâ, nitidiore; anfractibus 4½ planulatis, regulariter crescentibus, ultimo circa perforationem aper-tam excavato; aperturâ subrhombed; peristome simplici, tenui, margine columellari arcuatim reflexo.*

Diam. 1·80; altit. 1·00 poll.

Hab. Insula Burias.

**HELIX SAMARENSIS**, Pfr. *Hel. T. umbilicatâ, depresso-conoidea, tenui, obliquè striatulâ, fulvido-albâ, fasciis rufis ornatâ; basi planulatâ; suturâ linearî; anfractibus 4½ planiusculis, ultimo basi subcari-nato; umbilico angusto, pervio; aperturâ horizontali, ellipticâ; peristome simplici, margine supero expanso, basali latè reflexo, edentulo.*

Diam. 0·90; altit. 0·50 poll.

Hab. Insula Samar.

**HELIX BUTLERI**, Pfr. *Hel. T. imperforatâ, globosâ, tenui, lævi, apice obtuso albâ; spirâ semiglobosâ; suturâ mediocri; anfractibus 4½ planiusculis, ultimo inflato, pallidè lutescente, lineis par-vis confertis, crispulis vel interruptis ornatâ; columellâ subrectâ, latâ, profundiè intrante; aperturâ rotundato-lunatâ; peristome simplici, vix expanso.*

Diam. 1·25; altit. 1·00 poll.

Hab. Mountains of the Igorrotes.—Forma affinis *H. versicolori* Bornii.

**HELIX BECKIANA**, Pfr. *Hel. T. umbilicatâ, orbiculari, tenui, obliquè*

*striatâ, fuscâ, vix nitidulâ; spirâ parùm elatâ; suturâ linearî; anfractibus 6 planiusculis, ultimo deflexo, ad peripheriam angulato; angulo ad aperturam obsoleto; basi planiusculâ; umbilico mediocri, pervio; aperturâ ferè horizontali, subtrapezoidali; peristomate simplici, margine columellari brevi, basali reflexo, quasi in tuberculum incrassato.*

Diam. 0·85; altit. 0·40 poll.—Nueva Ecija.

**HELIX CUMINGII**, Pfr. *Hel. T. imperforatâ, depressâ, obtusè subcarinatâ, apice violaceo, obtusâ, obliquè striatâ, nigricanti-rufâ, epidermide rufâ, supernè maculis irregularibus, basi fasciis multis stramineo-cinereis ornata; anfractibus 4, supremis planiusculis, ultimo subinflato; columellâ rectè descendente, latè callosâ; aperturâ latâ, subquadrangulari; peristomate latè expanso, margine inferiore incrassato.*

Diam. 1·60; altit. 0·90 poll.

Hab. Insula Zebu.—Affinis *H. Zebuensi*, Brod.

**HELIX SCROBICULATA**, Pfr. *Hel. T. umbilicata, lenticulari, tenui, obliquè rugosâ, fulvido-albidâ, fusco-zonatâ, carinatâ; spirâ parùm elatâ, apice obtuso nitida, glabrâ; suturâ linearî; anfractibus 4½ convexiusculis, ultimo deflexo; carind subacutâ; umbilico angusto, pervio; aperturâ transversè pyriformi; peristomate simplici, continuo, supernè expanso, basi latè reflexo, unidentato; dente obtuso, extùs scrobiculum formante.*

Diam. 1·15; altit. 0·45 poll.

Hab. Insula Bohol.

Affinis *H. rotæ*, spirâ elatiore, carinâ simplici, subacutâ, et costis deficientibus diversâ.

**BULIMUS BREVICULUS**, Pfr. *Bul. T. imperforatâ, ovatâ, apice obtuso, obliquè obsoletè striatâ, nitidâ, albâ, epidermide lutescente deciduâ obductâ; anfractibus 6 angustis, convexiusculis; columellâ subrectâ, in laminam tenuem expansâ; aperturâ perobliquâ, transversè semiovali; peristomate simplici, expanso.*

Long. 1·15; diam. 0·75 poll.

Hab. Insula Romblon.

Affinis *Bulimo stabili*, Sow., formâ abbreviatâ, anfractibus convexiusculis et aperturâ diversis.

**BULIMUS CUMINGII**, Pfr. *Bul. T. imperforatâ, ovatâ, tenuissimâ, obliquè striatulâ, pellucidâ, albido-virente, ad suturam lineâ rufâ circumdatâ; anfractibus 4½ convexiusculis, ultimo obsoletè angulato, spiram paulò superante; columellâ planâ, subexcavatâ, fuscâ; aperturâ lunato-ovali; peristomate simplici, margine dextro latè expanso, basali subreflexo.*

Long. 1·35; diam. 0·95 poll.

Hab. Insula Camiguing.

**BULIMUS LIGNARIUS**, Pfr. *Bul. T. imperforatâ, conoideo-globosâ, solidâ, obliquè striatâ, subepidermide lignariâ nitidè nigricante, sursum pallescente, apice obtusiusculo, nitide fulvo; spirâ conoideâ;*

*anfractibus 5 convexis, ultimo spirā paulo breviore; columellā subrectā, extrorsū latē expansā, fusco-plumbeā; aperturā lunato-ovali, intūs nitidē cærulēscēte; peristomate simplici, reflexo, saturatē plumbeo, margine dextro valde arcuato.*

Long. 3·05; diam. 2·20 poll.

Hab. Provincia Cagayan insulæ Luçon.

α. *Ferè unicolor, epidermide pallidā, saturatius strigatā, fasciā unicā angustā, nigrā suprà, latiore albida infrà medium anfractus ultimi.*

β. *Epidermide albo-strigatā, fasciis pluribus angustis nigricantirufis, unicid latiore albida infrà medium anfractus ultimi.*

γ. *Minor, spira subelongata; color sicut in α.*

BULIMUS JUGLANS, Pfr. *Bul. T. imperforatā, elongato-globosā, apice obtuso, solidiusculā, obliquè distinctè striatā, unicolo, rufā; anfractibus 5 convexis, ultimo spiram æquante; columellā latā, albā, introrsū arcuatim prominente; aperturā magnā, irregulariter subovali, ad basin columellæ sinuatā, intūs pallidē lividā; peristomate vix reflexo, rufo-marginato.*

Long. 2·20; diam. 1·50 poll.

Hab. 'Mountains of Igorrotes.'

BULIMUS NYMPHA, Pfr. (*Achatina?*) *Bul. T. ovato-turritā, solidiusculā, lœvi, sulphureā, epidermide hydrophanā, ligned crebrè et latē strigatā, lineā suturali rufā et ared columellari nigricante ornatā; apice obtuso, nitide roseo; anfractibus 6 vix convexiusculis, ultimo  $\frac{1}{3}$  longitudinis æquante, obsoletè angulato; columellā rectā, planā, vix truncatulā; aperturā magnā, ovali, intūs albā; peristomate subsimplici, albo, margine dextro valde arcuato.*

Long. 2·00; diam. 0·95 poll.

Hab. Insula Luçon, 'San Miguel.'

Var. Testā rufā, sursūm pallescente, apice roseo; epidermide albida, peristomate rufo. Mt. Triga.

Differt a *Bulimo (Achatina) Boholensi* formâ ventricosiore, anfractu ultimo brevi, aperturā latā, peristomate non expanso.