

PROCEEDINGS


OF THE

ZOOLOGICAL SOCIETY

OF LONDON,

WITH ILLUSTRATIONS.

1849.



PRINTED FOR THE SOCIETY;  
SOLD AT THEIR HOUSE IN HANOVER SQUARE,  
AND BY MESSRS. LONGMAN, BROWN, GREEN, AND LONGMANS,  
PATERNOSTER ROW.

7. DESCRIPTION OF A NEW SPECIES OF CY THEREA.  
BY LOVELL REEVE, F.L.S., F.Z.S. ETC.

**CY THEREA NOBILIS.** *Cyth. testā orbiculari-cordatā, crassd, transversim concentricè liratā, liris rudibus, obtusis, subplanulatis, numerosis, crebris, valdē irregularibus, hic illic intermissis, non parallelis; lacte, epidermide tenui cornē, translucidā, induitā.*  
Long.  $4\frac{1}{2}$  in.; lat.  $2\frac{1}{2}$  in.; alt. 4 in.  
*Hab.* —?

This fine species, from the collection of A. L. Gubba, Esq., is distinguished by a peculiarity in the form and arrangement of the concentric ribs with which it is sculptured. They are very numerous, flattened, close-set, and extremely irregular, now narrow, now broad, each one varying irregularly in width and now and then suddenly intermittent. It is of a pure cream-colour, covered with a thin, horny, transparent epidermis. Mr. Gubba obtained it from a vessel in Havre-de-grace, but could not ascertain its locality.

8. DESCRIPTIONS OF TWENTY-FOUR NEW SPECIES OF HELICEA,  
FROM THE COLLECTION OF H. CUMING, ESQ.  
BY DR. L. PFEIFFER.

1. **STREPTAXIS GLABRA**, Pfr. *Str. testā umbilicatā, depressē ovatā, tenui, pellucidā, virescenti-albidā, omnino glabratā; spirā laterali, acutiusculā; suturā albo-marginatā; anfractibus 6 convexiusculis, penultimo inflato, ultimo antrorsum deviante; umbilico angusto, non pervio; aperturā perobliquā, semicirculāri, dente minuto parietis aperturalis munitā; peristomate albo, subincrastato, breviter reflexo.*

Diam. maj. 8, min. 6, alt. 5 mill.

*Hab.* Demerara.

2. **STREPTAXIS CUMINGIANA**, Pfr. *Str. testā perforatā, depresso-globosā, solidulā, glabratā, virenti-albidā; spirā sublaterali, conoidē; anfractibus 7 angustissimis, subplanis, penultimo prominulo, ultimo antrorsum deviante; aperturā perobliquā, auriformi, lamellā intrante parietis aperturalis coarctatā; peristomate simplice, breviter expanso, marginibus callo tenui junctis, dextro arcuato, infernè dente 1 acuto munito, basali stricto, dente 1 transversē elongato instructo, columellarī brevissimo, ad perforationem non perviam subreflexo.*

Diam. maj.  $6\frac{2}{3}$ , min.  $5\frac{1}{3}$ , alt. 4 mill.

Locality unknown.

3. **HELIX MONSSONI**, Pfr. *H. testā perforatā, turbinatā, tenui, levi, carinatā, striis incrementi et lineis confertissimis impressis, oblique antrorsum descendantibus subtilissimè decussatā, diaphanā, albida, rubro-unicingulatā; spirā conoidē, apice obtusiusculā; suturā submarginatā; anfractibus 6 subplanulatis, ultimo magno, infra cingulum carinato (carinā anticè ob-*

*soletd), basi convexiusculo; aperturd obliqud, subangulato-lunari; peristomate simplice, recto, margine columellari supra perforationem breviter reflexo.*

Diam. maj. 38, min. 33, alt. 23 mill.

Locality unknown.

4. **HELIX ALBICANS**, Pfr. *H. testd perforatd, depressd, striatuld, sublævigatd, nitidd, hyalino-albidd; spird vix elevatd; suturd impressd, marginatd; anfractibus 5 planiusculis, lentè accrescentibus, ultimo non descendente, subrotundato, circa perforationem impresso; aperturd verticali, latè lunari; peristomate simplice, acuto, margine columellari brevissimè reflexo.*

Diam. maj. 8, min. 7, alt. 4 mill.

Hab. in insulâ Jamaica.

5. **HELIX PHLOGOPHORA**, Pfr. *H. testd subperforatd, depresso-sulcata, tenuissimâ, striatuld, pellucidd, nitidd, fulvo-luted, flammulis angulatis et serrulatis rufis confertis pictd; spird parum elevatd, apice subpapillata; anfractibus  $3\frac{1}{2}$  convexis, rapidè accrescentibus, ultimo depresso, basi planiusculo; aperturd per obliqud, rotundato-lunari; peristomate simplice, recto, margine columellari subreflexo.*

Diam. maj.  $6\frac{1}{2}$ , min.  $5\frac{1}{2}$ , alt. 3 mill.

Locality unknown.

6. **HELIX SERICATULA**, Pfr. *H. testd perforatd, depressd, discoided, subtiliter et confertim costulatd, striatd, subsericd, griseo-corned, lineis brunneis irregularibus radiatd; spird planâ; anfractibus  $4\frac{1}{2}$  vix convexiusculis, ultimo subrotundato, juxta perforationem subimpresso; aperturd subverticali, latè lunari; peristomate simplice, recto, obtusiusculo, margine basali declivi, supernè reflexo, perforationem ferè occultante.*

Diam. maj.  $4\frac{1}{2}$ , min. 4, alt.  $2\frac{1}{3}$  mill.

Hab. ad Port Jackson (Mr. Strange).

7. **HELIX NOBILIS**, Pfr. *H. testd angustè umbilicatd, subturbinato-depressd, solidâ, striatd, lineis impressis concentricis et obliquis subtiliter decussatd, fulvd; spird parum elevatd, subturbinatd; anfractibus 6 parum convexis, ultimo medio circulo elevato, obtuso cincto, infra eum fasciâ saturatè castaneâ, deorsum dilutâ, ornato, circa umbilicum pallido; aperturd ampld, parum obliqud, latè lunari, intus margaritaceâ; peristomate simplice, recto, margine columellari ad umbilicum in laminam brevem, triangularem reflexo.*

Diam. maj. 53, min. 45, alt. 30 mill.

Hab. in insulâ Borneo, var. pallida in insulis Philippinis.

8. **HELIX BORNEENSIS**, Pfr. *H. testd obliquè perforatd, depressd, tenuiusculd, striis incrementi distinctis et lineis obliquis, impressis, crebris decussatd, saturatè fulvd; spird vix elevatd, obtusd; anfractibus 4 parum convexis, celeriter accrescen-*

*tibus, ultimo medio zonā nigricante, deorsum dilutā, ornato; suturā lined impressā marginatā; aperturā obliquā, amplā, transversē lunari-ovali, intus margaritacea, fascia pellucente; peristome simple, acuto, margine columellari in laminam brevem, triangularem, umbilicum semitegentem, reflexo.*

Diam. maj. 52, min. 42, alt. 25 mill.

Hab. in insulā Borneo.

9. *HELIX AFRICANA*, Pfr. *H. testā perforatā, depressā, tenui, nitidā, minutissimè striatula, lineis confertis, concentricis, impressis sub lente minutissimè decussatā, rufo-fuscā vel pallidè corned; spirā brevissimè conoidea, apice subelevata; suturā submarginatā; anfractibus 7 vix convexiusculis, sensim accrescentibus, ultimo carinato (carinā anticè obsoletā), non descendente, basi paulo convexiore; aperturā depressā, latā, lunari; peristome simple, recto, acuto, margine columellari supra perforationem brevissimè reflexo.*

Diam. maj. 26, min. 23, alt. 13 mill.

Hab. ad Axim in littore occidentalī Africæ.

10. *HELIX SANDVICENSIS*, Pfr. *H. testā umbilicatā, discoidē, striatā, nitidulā, luteo-corned; spirā plana; suturā impressā; anfractibus 5–6 lentè accrescentibus, ultimo depresso, basi vix convexiore; umbilico lato, dimidium ferè diametri occupante; aperturā parum obliquā, lunari-rotundatā; peristome simple, recto, tenui, marginibus conniventibus.*

Diam. maj. 18, min. 15, alt. 5 mill.

Hab. in insulis Sandwich.

11. *HELIX JACQUINOTI*, Pfr. *H. testā umbilicatā, fornicate-conoidea, solidulā, acutē carinatā, confertim arcuato-costatā, albo et fusco variegatā; spirā conoidea, obtusa; anfractibus 8 angustis, omnibus carinatis (carinā exsertā, compressā, costis decurrentibus denticulatā), ultimo basi vix convexiusculo, radiatim striato; umbilico extus laminā horizontali coarctato, intus lato; aperturā depressā, securiformi, lamellis 6 intrantibus munita: 2 in pariete aperturali elongatis, 1 columellari et 3 in margine basali profundis, vix conspicuis; peristome simple, recto, acuto, margine basali in lamellam umbilici introitum circumclaudentem continuato.*

Diam. maj. 9, min.  $8\frac{1}{2}$ , alt. 5 mill.

Hab. in insulā Tahiti, et in insulis Marquesas.

12. *HELIX COARCTATA*, Pfr. *H. testā umbilicatā, depressā, distanter arcuato-costatā et sub lente minutissimè spiraliter striatā, fuscula, brunneo-tessellatā; spirā fornicate, supernè depressā; anfractibus  $8\frac{1}{2}$  angustis, carinatis, ultimo infra penultimum recedente, tertiam paginā inferā penultimi partem liberam relinquente, basi vix convexiusculo, obsoletè radiatim costato, distinctius concentricè striato; umbilico lato, extus laminā horizontali coarctato; aperturā depressā, securiformi,*

*lamellis 6 intrantibus munita: 2 in pariete aperturali, 2 in margine basali, 1 in supero, 1 dentiformi in columella; peristome simple, recto, acuto, margine basali retrorsum in lamina, umbilicum coarctantem, continuata.*

Diam.  $6\frac{1}{3}$ , alt.  $3\frac{1}{2}$  mill.

Hab. in insulâ Tahiti.

13. **HELIX NYMPHA**, Pfr. *H. testa imperforata, globoso-depressa, tenui, obliquè striatula, nitida, diaphana, virenti-albida; spiræ brevissimæ, apice obtusa; suturæ albo-flosæ; anfractibus 4 subplanis, rapidè accrescentibus, ultimo subdepresso, basi convexo; columella intrante, subverticali, compressa, albæ; apertura obliqua, latè lunari; peristomate simple, tenui, castaneo-limbato, margine supero recto, basali breviter reflexo, cum columella angulum obtusum formante.*

Diam. maj. 32, min. 26, alt. 18 mill.

Hab. in insulis Philippinis.

14. **HELIX TRICOLOR**, Pfr. *H. testa lenticulari-conoidæ, tenui, carinata, undique confertim concentricè striata, hyalina, supernè lineis albæ et ad suturam fascia alba, castaneo-punctatæ ornata; spiræ brevi, conoidæ, apice obtusa; anfractibus 4 planiusculis, ultimo carinæ alba, lineis castaneis marginata et articulata munito, supernè et ad carinam subito deflexo, basi juxta columellam subgibbo; apertura angusta, perobliqua, subquadrangulari; peristomate simple, castaneo-limbato, margine dextro recto, basali breviter reflexo, columellari perdeclivi, introrsum dilatato, excavato, saturatè castaneo.*

Diam. maj. 34, min. 29, alt. 17 mill.

Hab. St. Christoval, ins. Salomonis.

15. **HELIX RECEDENS**, Pfr. *H. testa imperforata, subsemiglobosa, solidæ, carinata, supernè confertim costulato-striata, pallide carnea; spiræ fornicate; suturæ vix impressæ; anfractibus 6 planiusculis, lentè accrescentibus, ultimo a medio infra penultimum recedente, basi planiusculo, striato; carinæ rufolineata; apertura obliqua, angulato-lunari; peristomate sub-simple, margine dextro recto, basali subincrassato, columellari brevissimè reflexo.*

Diam. 12, alt. 7 mill.

Locality unknown.

16. **HELIX SALLEANA**, Pfr. *H. testa imperforata, conicæ, tenuiscula, striatula et impressionibus obsoletis rugosula, parum nitida, diaphana, cinereo-lutescente, ad peripheriam fasciis 2 fusco-viridibus, punctisque castaneis ornata; spiræ conicæ, acutiscula; anfractibus 5 convexiusculis, ultimo lineis concentricis impressis notato, subangulato, basi parum convexo; apertura parum obliqua, lunari-ovali, intus nitida, concolore, fasciis nigricantibus; peristomate tenui, rectangularè expanso et reflexiusculo, albo, margine columellari supernè dilatato, calloso.*

Diam. maj. 30, min. 24, alt. 24 mill.

*Hab.* in ripis fluvii St. Johan. Guatimalæ (Sallé).

17. **HELIX PLATYSTYLA**, Pfr. *H. testā imperforatā, conicā, solidā, costulato-striatā, albida, lined rufā ad suturam ornatā; spira conicā, acutiuscula; anfractibus 6 convexiusculis, sensim acresentibus, ultimo obsoletē angulato, basi subplano; aperturā obliquā, latē lunari; peristomate simplice, marginibus subparallelis, dextro breviter expanso, columellari supernè perditato, calloso.*

Diam. maj. 22, min. 19, alt. 19 mill.

*Hab.* in insulis Moluccis?

18. **HELIX BREVIPILA**, Pfr. *H. testā umbilicatā, globoso-depressā, pilis brévissimis, rigidis, quincuncialiter dispositis asperā, haud nitidā, saturatē brunneā; spirā parum elevatā, obtusā; anfractibus vix 5 convexiusculis, ultimo rotundato, anticē subdeflexo, circa umbilicum angustum subcompressō; aperturā obliquā, rotundato-lunari, intus nitidā; peristomate tenui, brunneo-carneo, breviter expanso, marginibus conniventibus, columellari subdilatato-reflexo.*

Diam. maj. 12, min. 10, alt. 6½ mill.

*Hab.* in orā orientali Novae Hollandiæ (Mr. Strange).

19. **HELIX BASKERVILLEI**, Pfr. *H. testā umbilicatā, globoso-depressā, solidā, striis incrementi rugulosis, lineisque spiralibus impressis subgranulatā, olivaceo-fuscā, parum nitidā; spirā subconoideo-elevatā, apice obtusiuscula; suturā impressā, crenulatā; anfractibus 6½ angustis, convexiusculis, ultimo rotundato, anticē vix descendente; aperturā parum obliquā, lunari, dente linguiformi albo parietis aperturalis, obliquè intrante, coarctatā; peristomate validē carneo-labiato, margine dextro arcuato et basali subhorizontali, latē subdentato latē expansis et reflexis, columellari brevi, perdilatato, umbilicum angustum semitegente.*

Diam. maj. 24, min. 19, alt. 14 mill.

*Hab.* Vancouver's Island (Lieut. Baskerville).

20. **HELIX CONNIVENS**, Pfr. *H. testā angustè umbilicatā, depresso, solidā, striis incrementi distinctis, lineisque subtilibus concentricis sub lente decussatā, corneo-stramineo, nitidulā; spirā parum elevatā; suturā impressā; anfractibus 6 parum convexis, ultimo anticē non descendente, peripheria subangulata, basi convexiore; aperturā obliquā, latē lunari, intus albida; peristomate intus valide albo-labiato, marginibus conniventibus, dextro acuto, parum expanso, basali breviter reflexo, columellari supernè dilatato, patente.*

Diam. maj. 26, min. 22, alt. 14 mill.

*Hab.* Liew-Kiew.

21. **HELIX GALACTOSTOMA**, Pfr. *H. testā umbilicatā, convexo-*

*orbiculatā, solidā, striatā, sub lentē granulatā, fulvā; spira brevi, fornicatā, obtusā; anfractibus  $4\frac{1}{2}$  convexiusculis, sensim accrescentibus, penultimo angulato, ultimo subdepresso, anticē deflexo, basi subangulatim in umbilicum, mediocrem, pervium,  $\frac{1}{6}$  diametri vix superantem descendente; aperturā perobliquā, lunari-ovali, intus lacteū; peristomate simplice, fusculo-limbato, marginibus conniventibus, callo junctis, dextro brevissimè expanso, basali subreflexo, columellari albo, supra umbilicum dilatato-reflexo.*

Diam. maj. 36, min. 31, alt. 18 mill.

Hab. in insulā Madagascar.

22. **HELIX ROSARIUM**, Pfr. *H. testā umbilicatā, depressā, tenui, supernē subtilissimè granulatā, diaphanā, pallide fulvā, flammis a suturd exaeuntibus, cingulisque 3 interruptis, moniliformibus rufis ornatā; spira planā; anfractibus vix 5 convexis, ultimo subdepresso, anticē non descendente, basi radiatim striatulo et lineis impressis spiralibus distantibus notato, circa umbilicum mediocrem, infundibuliformem subcompresso; aperturā parum obliquā, lunato-subtriangulari; peristomate albo-labiato, breviter reflexo, margine supero ab insertione primum ascendentē, tum sinuato, basali strictiusculo, repando, columellari brevi, triangulatim patente.*

Diam. maj. 21, min. 17, alt. 9 mill.

Locality unknown.

23. **BULIMUS (PARTULA) DECUSSATULUS**, Pfr. *B. testā perforatā, orato-conicā, tenui, striis incrementi lineisque spiralibus minutissimè decussatula, vix nitidula, fulvescenti-albidā, diaphanā; spira brevi, conicā, obtusiuscula; suturd mediocri; anfractibus  $4\frac{1}{2}$  convexis, ultimo  $\frac{5}{9}$  longitudinis equante, rotundatā; columella subplicatā, recedente; aperturā angulato-ovali; peristomate simplice, tenui, marginibus conniventibus, dextro campanulatim expanso, columellari super perforationem reflexo.*

Long. 15, diam.  $8\frac{2}{3}$  mill.; ap. 9 mill. longa,  $6\frac{1}{2}$  lata.

Hab. in insulā Navigatorum.

24. **BULIMUS (PARTULA) NAVIGATORIUS**, Pfr. *B. testā dextorsd, perforata, oblongo-ovatā, solidā, obsoletē granulato-striatā, nitidula, fulvā, lineis confertis saturationibus signatā; spira conicā, acutiusculā; suturd levi, albo-marginatā; anfractibus 5 planiusculis, ultimo spiram superante; aperturā oblongā, angustā, intus albida, dente calloso parrulo profundo in ventre anfractū penultiimi munitā; peristomate subincrasato, intus valide albo-labiato, marginibus parallelis, dextro breviter expanso, medio subdentato, columellari dilatato, plano, reflexo.*

Long. 23, diam. 11 mill.; ap. (c. perist.) 13 mill. longa, 8 lata.

Hab. in insulā Navigatorum.