

PROCEEDINGS  
OF THE  
ZOOLOGICAL SOCIETY  
OF LONDON.



---

PART XXII.

1854.

---

PRINTED FOR THE SOCIETY ;  
SOLD AT THEIR HOUSE IN HANOVER SQUARE,  
AND BY MESSRS. LONGMAN, BROWN, GREEN, AND LONGMANS,  
PATERNOSTER ROW.

*intus plicis 6-7 brevibus, albis munito, columellari calloso, subdilatato.*

Long. 10, diam. 5 mill.

Hab. New Zealand.

4. **MELAMPUS SPIRALIS**, Pfr. *M. testa subrimata, biconica, solidula, striatula, nitidula, alba; spira conoidea, acute mucronata; sutura linearis; anfr. 10, superis planis, distincte spiraliter striatis, ultimo  $\frac{3}{4}$  longitudinis fere æquante, spiraliter obsoletissime striato, infra suturam excavato, tum obtuse angulato, deorsum regulariter attenuato, basi compresso; apertura subverticali, oblonga, basi rotundata; plicis parietalibus sub 2, superiore lamellæformi, intrante, altera immediate infra illam, obsoleta; plica columellari alta, extrorsum subhorizontaliter porrecta; perist. acuto, margine dextro superne angulato, intus plicis transversis 7-8 munito, columellari parum dilatato, subpatente.*

Long. 16, diam. max. 9 mill.

Hab. —?

5. **MELAMPUS CRISTATUS**, Pfr. *M. testa imperforata, oblongo-ovata, solida, spiraliter confertim striata, sericea, luteo-fulva; spira brevi, convexa, mucronata; sutura canaliculata; anfr. 8, convexiusculis, penultimo latiore, ultimo  $\frac{2}{3}$  longitudinis formante, supra medium ventroso, basi attenuato; apertura vix obliqua, perangusta, basi bisinuosa; plicis parietalibus 3, suprema nodiformi, 2 subtransversis, extus in cristam callosam, spiralem continuatis; plica columellari valida, fere verticali; perist. acuto, margine dextro intus callo crasso, albo, confertim transverse plicato munito, columellari subdilatato, appresso.*

Long.  $9\frac{1}{2}$ , diam. 6 mill.

Hab. Philippine Islands.

6. **PEDIPES TRIDENS**, Pfr. *P. testa imperforata, globoso-conica, tenui, obsolete et confertim spiraliter striata, saturate fusca; spira turbinata, acutiuscula; sutura profunda, marginata; anfr. 5, convexis, ultimo  $\frac{3}{5}$  longitudinis formante, inflato; apertura fere diagonalis, subsemicirculari; plica parietali mediocri; labio columellari calloso, dentibus 2 distantibus munito; perist. simplice, acuto, edentulo.*

Long.  $3\frac{3}{4}$ , diam. max.  $2\frac{1}{2}$  mill.

Hab. Bermuda.

## 2. DESCRIPTIONS OF SIXTEEN NEW SPECIES OF HELICEA, FROM THE COLLECTION OF H. CUMING, ESQ.

BY DR. L. PFEIFFER.

1. **VITRINA AUREA**, Pfr. *V. testa ambitu subauriformi, depressa, nitidissima, pellucida, aurea; spira plana, vertice vix prominulo; sutura levissima, marginata; anfr. 3 subplanis, rapide accrescentibus, ultimo depresso, peripheria obsoletissime subangulato,*

*basi vix convxiore, lato; apertura perobliqua, ovali, intus submargaritacea; perist. simplice, margine dextro antrorum subdilatato, columellari leviter inflexo.*

Diam. maj. 13, min. 9, alt.  $4\frac{2}{3}$  mill.

Hab. Salomon's Islands.

2. **SUCCINEA SIMPLEX**, Pfr. *S. testa ovata, tenui, striata, pellucida, nitida, succinea; spira brevi, conica, vertice submammillato, acutiusculo; anfr.  $2\frac{1}{2}$  convexis, ultimo  $\frac{2}{3}$  longitudinis æquante; columella substricta recedente, leviter callosa; apertura obliqua, incumbente, angulato-ovali; perist. simplice, recto, marginibus regulariter conniventibus.*

Long.  $9\frac{1}{2}$ , diam. 6, alt.  $4\frac{1}{2}$  mill.

Hab. Salomon's Islands.

3. **SUCCINEA REPANDA**, Pfr. *S. testa ovato-oblonga, tenui, confertissime capillaceo-striata, vix sericea, pallide griseo-cornea; spira conica, acuta, plerunque erosa; anfr.  $3\frac{1}{2}$ , penultimo convexiusculo, ultimo basi repando; columella superne subplicata, callosa, leviter arcuata; apertura obliqua, angulato-ovali, intus submargaritacea; perist. simplice, recto, marginibus callo tenuissimo junctis.*

Long. 12, diam.  $6\frac{1}{2}$ , alt. 5 mill.

Hab. Banks of the river Negro, New Granada.

4. **HELIX GENERALIS**, Pfr. *H. testa imperforata, globoso-turbinata, striatula et rugulis obliquis præcipue subtus sculpta, non nitida, sub epidermide luteo-virente alba; spira convexo-conoidea, obtusula; sutura levi, submarginata; anfr.  $4\frac{1}{2}$ , convexiusculis, celeriter accrescentibus, ultimo antice vix descendente, supra peripheriam lira plana latiuscula cincto, basi parum convexo; columella vix obliqua, compressa, excavata, alba; apertura diagonali, oblique truncato-ovali, intus lactea; perist. tenui, marginibus conniventibus, dextro vix expansiusculo, basali incrassato, breviter reflexo, cum columella angulum obtusum formante.*

Diam. maj. 57, min. 45, alt. 40 mill.

Hab. Philippine Islands.

5. **BULIMUS EXCORIATUS**, Pfr. *B. testa aperte umbilicata, ovato-acuminata, tenuiuscula, striatula, sub lente punctulata, non nitente, carne; spira conica, acuta; anfr. 6, convexiusculis, ultimo spiram superante, subinflato; apertura parum obliqua, acuminate-ovali; perist. tenui, expanso, marginibus fere contiguis, callo tenui junctis, dextro medio dilatato, columellari perdilatato, patente.*

Long. 39, diam. 19 mill.

Hab. Andes of Peru (*Captain Keppell*).

6. **BULIMUS CRETACEUS**, Pfr. *B. testa umbilicata, ovato-conica, solidula, striatula et impresso-punctata, opaca, cretacea; spira*

*convexo-conica, apice acutiuscula ; anfr. 6, vix convexiusculis, ultimo spiram paulo superante, antice subascendente, basi juxta umbilicum profundum, subcompressum rotundato ; apertura vix obliqua, oblongo-ovali, intus fusca ; perist. tenui, albo-limbato, marginibus convergentibus, callo junctis, dextro late expanso, columellari fusco, dilatato, fornicatim patente.*

Long. 36, diam. 18 mill.

*Hab.* Eastern Islands (*Captain Keppell*).

**7. BULIMUS TRISTIS**, Pfr. *B. testa compresse umbilicata, oblongo-conica, tenui, laevigata, nitidula, albida, fasciis et strigis vel seriebus macularum quadratarum, violaceo-fuscarum ornata ; spira convexo-conica, apice acuta ; anfr. 6 $\frac{1}{2}$ , vix convexiusculis, ultimo spira paulo breviore, basi angustato, rotundato ; columella leviter arcuata ; apertura vix obliqua, ovali ; perist. tenui, margine dextro breviter expanso, columellari luteo vel albo, superne dilatato, fornicatim reflexo.*

Long. 28, diam. 11 mill.

*Hab.* New Granada.

**8. BULIMUS HISTRIO**, Pfr. *B. testa subperforata, ovato-conica, tenui, laevigata, nitida, lutescenti-alba, serie macularum castanearum, subquadratarum supra suturam picta ; spira conica, apice nigra, obtusula ; anfr. 5 $\frac{1}{2}$ , convexiusculis, ultimo spiram vix superante, infra medium nigro-fusco bizonato, basi rotundato ; apertura parum obliqua, elliptico-ovali ; perist. simplece, recto, margine dextro arcuato, columellari subverticali, superne late reflexo, perforationem fere tegente.*

Long. 13, diam. 7 $\frac{1}{2}$  mill.

*Hab.* Tanna, New Hebrides.

**9. BULIMUS COLUMBIENSIS**, Pfr. *B. testa breviter rimata, subperforata, oblongo-turrita, solida, sublaevigata, parum nitida, flavidio-albicante ; spira elongata, subconvexo-turrita, apice cornea, acutiuscula ; anfr. 7, vix convexiusculis, ultimo  $\frac{3}{7}$  longitudinis æquante, basi subpresso ; columella stricta, rececente ; apertura parum obliqua, oblongo-ovali ; perist. recto, acuto, margine dextro leviter arcuato, columellari subito reflexo, patente.*

Long. 28, diam. 11 $\frac{1}{2}$  mill.

*Hab.* Columbia.

**10. BULIMUS SIMPLICULUS**, Pfr. *B. testa perforata, conico-ovata, solida, ruguloso-striata (lineis spiralibus sub lente obsolete decussata), nitidula, alba, punctis pellucidis passim conspersa ; spira convexo-conica, apice cornea, obtusula ; sutura minute crenulata ; anfr. 6, modice convexis, ultimo spira paulo breviore, basi subrotundato ; columella substricta, superne subplicata ; apertura vix obliqua, oblongo-ovali ; perist. simplece,*

*recto, margine dextro leviter arcuato, columellari dilatato, fornicate reflexo.*

Long. 19½, diam. 9½ mill.

Hab. —?

11. **BULIMUS MARMATENSIS**, Pfr. *B. testa perforata, globosoconica, tenui, striatula, pellucida, pallide griseo-fulva; spira conica, obtusiuscula; anfr. 4½, modice convexis, ultimo spiram vix superante, rotundato; columella leviter arcuata; apertura parum obliqua, rotundato-ovali; perist. simple, recto, margine dextro perarcuato, columellari superne dilatato, supra perforationem angustam fornicate reflexo.*

Long. 17, diam. 10 mill.

Hab. Marmato, New Granada.

12. **BULIMUS HAPLOCHROUS**, Pfr. *B. testa perforata, ovato-fusiformi, tenui, sublævigata, non nitente, diaphana, corneo-albida; spira elongata, convexo-turrita, acutiuscula; anfr. 7, vix convexiusculis, ultimo spira paulo breviore, basi attenuato; columella leviter arcuata; apertura obliqua, subelliptica, basi subangulata; perist. simple, recto, margine dextro regulariter arcuato, columellari sursum dilatato, fornicate-reflexo.*

Long. 30, diam. 12 mill.

Hab. —?

13. **BULIMUS DIAPHANUS**, Pfr. *B. testa anguste perforata, turrito-oblonga, tenui, striatula, haud nitente, diaphana, epidermide rufa, transversim lirata induita; spira elongato-conica, apice obtusula; anfr. 6, convexiusculis, ultimo ¾ longitudinis subæquante, basi rotundato; columella substricta; apertura vix obliqua, oblongo-ovali; perist. simple, acuto, margine dextro recto, leviter arcuato, columellari vitreo, superne fornicate reflexo.*

Long. 15, diam. 7 mill.

Hab. S. Thomas, W. Indies (*Bland*).

14. **PARTULA PACIFICA**, Pfr. *P. testa compresse umbilicata, ovato-conica, tenuiuscula, sublente minutissime spiraliter striata, nitidula, straminea, lineis albis et fulvis radiata; spira conica, apice acuta; sutura profunda, submarginata; anfr. 5½, convexis, ultimo spira paulo breviore, basi subattenuato; columella superne recedente, fornicate; apertura vix obliqua, truncato-oblonga; perist. albo, expanso, margine dextro superne arcuato, columellari late patente et reflexo.*

Long. 22, diam. 10 mill.

Hab. Pacific Islands.

15. **ACHATINA VIRENS**, Pfr. *A. testa cylindraceo-turrita, tenuissima, leviter striatula (striis ad suturam distinctioribus), nitidissima, pellucida, virenti-cornea; spira elongata, superne sensim attenuata, apice obtusula; sutura subcrenulata; anfr.*

8, convexiusculis, ultimo  $\frac{1}{3}$  longitudinis vix æquante, convexiore; columella perarcuata, abrupte truncata; apertura verticali, irregulariter ovali; perist. simplice, recto, margine dextro leviter arcuato.

Long. 18, diam. 8 mill.

Hab. —?

16. ACHATINA MANIENSIS, Pfr. *A. testa subrimata, ovato-oblonga, solidula, sublævigata, pallide cornea; spira convexoturrita, apice obtusa; sutura marginata; anfr. 6, vix convexiusculis, ultimo  $\frac{2}{5}$  longitudinis subæquante, basi subcompresso; columella verticali, introrsum subtorta, ad basin aperturæ sub-rhombeo-ovalis truncatula; perist. simplice, obtuso, margine columellari reflexiusculo, adnato.*

Long. 7, diam. 3 mill.

Hab. Mani, Sandwich Islands (Dr. Newcomb).

### 3. DESCRIPTIONS OF ELEVEN NEW SPECIES OF CYCLOSTOMACEA, FROM THE COLLECTION OF H. CUMING, ESQ.

BY DR. L. PFEIFFER.

1. CYCLOSTOMA (CYCLOPHORUS) ALABASTRINUM, Pfr. *C. testa umbilicata, subdepresso-turbinata, solida, sublævigata, spiraliter minutissime striata, alabastrina; spira turbinata, sursum livido-fusca, apice convexiuscula; anfr. 5, convexis, celeriter accrescentibus, ultimo peripheria obsoletissime subangulato et antice rudimenta fasciæ castaneæ exhibente, basi inflato; umbilico angusto, vix pervio; apertura vix obliqua, circulari; perist. fulvo-carneo, continuo, breviter adnato, crasso, undique reflexo, margine sinistro subdilatato.*

Diam. maj. 45, min. 34, alt. 25 mill.

Hab. Ceylon (Mr. Thwaites).

2. CYCLOSTOMA (CYCLOPHORUS) POLYNEMA, Pfr. *C. testa umbilicata, depresso-turbinata, tenuiuscula, oblique striatula et liris filiformibus confertissimis, subtus obsoletioribus, cincta, superne flammis angulosis lœte castaneis et albidis variegata; spira brevi, turbinata, apice obtusula; anfr.  $4\frac{1}{2}$ , convexiusculis, rapide accrescentibus, ultimo ad suturam antice subplanato, peripheria obsoletissime angulato et fascia lata, nigro-fusca, deorsum diluta ornato, circa umbilicum mediocrem pallido; apertura obliqua, subrotundata, latiore quam alta; perist. albo, continuo, breviter adnato, margine dextro expansiusculo, columellari breviter revoluto.*

Diam. maj. 33, min. 26, alt. 17 mill.

Hab. —?

3. CYCLOSTOMA (CYCLOPHORUS) EXCELLENS, Pfr. *C. testa umbilicata, subturbinato-depressa, solida, confertim spiraliter*