

PROCEEDINGS
OF THE
ZOOLOGICAL SOCIETY
OF LONDON.



PART XIV.

1846.

PRINTED FOR THE SOCIETY,
BY R. AND J. E. TAYLOR, RED LION COURT, FLEET STREET.

19. *BULIMUS NILAGIRICUS*, Pfr. *Bul. testá rimato-perforatá, oblongo-turritá, solidá, opacá, lineis impressis confertissimis subundulatis obsolete sculptá, fuscá, albido obliquè strigatá; spirá regulariter turritá, apice obtusiusculo; anfractibus 8 vix convexiusculis, ultimo $\frac{1}{3}$ longitudinis subæquante, basi subcompresso; aperturá ovali; peristomate expanso, latè albo-labiato, margine dextro supernè subangulato, columellari usque ad basin dilatato, patente.*

Long. $28\frac{1}{2}$, diam. 8 mill.

From the Neelgherries, East Indies.

20. *BULIMUS ZONULATUS*, Pfr. *Bul. testá perforatá, oblongo-conicá, tenui, lævigatá, opacá, pallidè stramineá, seriebus 2 macularum fuscarum pellucidarum cinctá, basi lineis 2 castaneis ornatá; spirá conicá, acutiusculá, apice corneo; anfractibus 6 convexiusculis, ultimo spiram vix superante; columellá strictá; aperturá ovali-oblongá; peristomate acuto, tenui, margine columellari a basi dilatato, membranaceo, angulatim latè reflexo, perforationem ferè occultante.*

Long. 18, diam. 9 mill.

From Cabanatuan, province of Nueva Ecija, island of Luzon; found by Mr. H. Cuming.

“Description of nine new species of *Helicea*, collected by H. Cuming, Esq.,” by Dr. L. Pfeiffer:—

1. *HELIX LUCIDELLA*, Pfr. *Hel. testá minutá, perforatá, depressá, striatuld, nitidissimá, brunneá; spirá subplanulatá; suturá albomarginatá; anfractibus 4 planis, ultimo basi vix convexiore, medio impresso, angustissimè perforato; aperturá obliquè lunari; peristomate simplice, obtuso, margine columellari declivi, vix incrassato.*

Diam. $3\frac{2}{3}$, alt. 2 mill.

Found on the island of Luzon.

2. *HELIX ARCTISPIRA*, Pfr. *Hel. testá umbilicatá, depressá, subdiscoideá, confertim costatá, albidd, epidermide tenui fuscescente indutá; spirá convexiusculá; anfractibus $5\frac{1}{2}$ convexis, angustissimis; umbilico lato, perspectivo; aperturá parvuld, subverticali, lunato-orbiculari; peristomate simplice, acutá.*

Diam. $2\frac{1}{2}$, alt. $1\frac{1}{4}$ mill.

From the island of Juan Fernandez.

Intermediate between *H. epidermia*, Aut., and *tessellata*, Mühlf.

3. *HELIX CYATHELLUS*, Pfr. *Hel. testá umbilicatá, conicá, obliquè costatá, tenuiusculá, unicolore cornéá; spirá pyramidatá, acutiusculá; anfractibus 9 angustissimis, cariná filiformi cinctis, ultimo basi planiusculo, sublævigato; umbilico majusculo, pervio; aperturá depressá, angulato-lunari; peristomate simplice, margine supero brevi, recto, basali leviter arcuato, brevissimè reflexo.*

Diam. $5\frac{1}{3}$, alt. 4 mill.

From the island of Panay.

4. *HELIX DOLIOLUM*, Pfr. *Hel. testá perforatá, turbinatá, confer-*

tissimè et minutè costulato-striatá, pellucidá, non nitente, cornedá; spirá turbinatá, apice obtusiusculo; anfractibus 5 convexiusculis, ultimo basi subplanato; aperturá depressá, latè lunari; peristomate simplicissimo, recto.

Diam. $3\frac{1}{2}$, alt. $2\frac{1}{2}$ mill.

From Sibonga, island of Zebu.

5. *BULIMUS DILATATUS*, Pfr. *Bul. testá imperforatá, ovato-conicá, obtusiusculá, solidá, obliquè striatá, subtilissimè punctatá, castaned, supernè fulvá; anfractibus 6 planiusculis, ultimo ad suturam albo-unifasciato, spirá multò breviorè; columellá subrectá, callosá, albá, dilatatá; aperturá obliqua, latè semiovali, intus lacteá; peristomate subincrassato, expanso, margine busuli reflexo, appresso.*

Long. 34, diam. 22 mill.

Island of Luzon.

6. *BULIMUS ELONGATULUS*, Pfr. *Bul. testá imperforatá, subulatá, solidulá, sub epidermide tenuissimè striatá (interdum obsolete decussatá), albá; spirá subulatá, acutá; anfractibus 8 planiusculis, ultimo tertiam longitudinis partem ferè aequante; columellá breviter recedente, callosá, planá; aperturá oblongo-ovuli; peristomate simplice, margine dextro antrorsum subarcuato, columellari subincrassato, appresso.*

Long. 24, diam. $5\frac{1}{2}$ mill.

Island of Luzon.

7. *BULIMUS GRATELOUPI*, Pfr. *Bul. testá imperforatá, ovato-oblongá, ruguloso-striatá, tenuissimá, nitidá, pellucidá, stramineo-ulbidá; anfractibus 6-7 convexiusculis, ultimo spirá paulò breviorè; columellá callosá, retrorsum flexuosá; aperturá semiovali; peristomate simplice, acuto, margine dextro antrorsum arcuato.*

Long. 18, diam. 8 mill.

From the islands of Luzon and Panay.

8. *BULIMUS PHILIPPINENSIS*, Pfr. *Bul. testá imperforatá, ovato-turbinatá, solidá, nigricante, strigis obliquis epidermidis hydrophanæ griseo-fuscæ ornatá; spirá conicá, obtusiusculá, nudá, pulidá; anfractibus 6 convexis, diametro celeriter accrescentibus, ultimo spirá paulò breviorè; columellá vix obliquá, subtortá, carnéa; aperturá lunato-orbiculari, intus lacteá; peristomate subincrassato, breviter reflexo, nigro-limbato, margine dextro valdè arcuato, columellari dilatato, expanso.*

β. *Testa epidermide fusca, saturatius strigata ferè omninò obducta, fasciis variis nigricantibus circumdata.*

Long. 63, diam. 41 mill.

From the islands of Luzon and Marinduque.

9. *HELIX REEVEANA*, Pfr. *Hel. testá umbilicatá, subdiscoicá, tenuiusculá, obliquè striatá, albidá, zonis 3-5 rufis ornatá; spirá vix convexiusculá, obtusá; anfractibus $4\frac{1}{2}$ -5 planiusculis, ultimo anticè subitò deflexo, basi concentricè et confertim striato; umbilico me-*

diocri, pervio ; aperturâ subhorizontali, transversè ovali ; peristomate subsimplice, reflexo, marginibus junctis.

Diam. 30, alt. 12 mill.

Island of Zebu.

This shell has been often mistaken for *H. Lasallii*, Eydoux, and is in many collections under that name ; but *H. Lasallii* is not this species. It is quite congruent with *H. meretria*, Sow.

April 28, 1846.

No business was transacted.

May 12, 1846.

Richard C. Griffith, Esq., in the Chair.

Mr. H. E. Strickland exhibited a species of *Corvus*, discovered by Capt. H. M. Drummond, 42nd R. H., which the latter gentleman proposes to name *Corvus collaris*. In size and form it is closely allied to the Common Jackdaw, *Corvus monedula*, but differs in the much lighter silvery grey of the cheeks, occiput and nape, which passes into a well-marked patch of pure white on each side of the neck. The black on the crown is of less extent than in *Corvus monedula*, and the lower parts are of a slaty grey.

Capt. Drummond states that in Macedonia and Thessaly this bird takes the place of *C. monedula*, which is common in the south of Greece, and does not there differ from the Jackdaw of Britain.

May 26, 1846.

No business was transacted.