

# PROCEEDINGS

OF THE

# ZOOLOGICAL SOCIETY

OF LONDON.



---

PART XIV.

1846.

---

PRINTED FOR THE SOCIETY,

BY R. AND J. E. TAYLOR, RED LION COURT, FLEET STREET.

December 8, 1846.

George Gulliver, Esq., F.R.S., in the Chair.

A paper was read containing descriptions of 38 new species of Land-shells, in the collection of Hugh Cuming, Esq., by Dr. L. Pfeiffer :—

1. **PARMACELLA CUMINGI**, Pfr. *Parm. testā depresso-semiovatā, tenuissimā, striatā, lineis spiralibus subtiliter decussatā, diaphanā, pallidē virenti-corned vel (in adultis) succined; spirā vix prominulā, subpapillatā; anfractibus 2; columellā arcuatā, acutā.*

Long. 6, lat. 6, alt.  $2\frac{3}{4}$  mill.

From the island of Juan Fernandez (H. Cuming).

2. **SUCCINEA PALLIDA**, Pfr. *Succ. testā ovato-conicā, solidiusculā, longitudinaliter ruguloso-striatā, diaphanā, pallidē stramineā; spirā acutā; anfractibus 4 convexis; columellā leviter arcuatā, filari, supra basin aperturā ovalis subtruncatā; peristomate margine subincrassato.*

Long. 13, lat. 7, alt.  $6\frac{1}{2}$  mill.; apert.  $8\frac{1}{2}$  mill. longa.

From Tahiti (H. Cuming).

3. **SUCCINEA SEMIGLOBOSA**, Pfr. *Succ. testā ovato-semiglobosā, tenui, lēvigatā, nitidissimā, lutescenti-corned; spirā vix prominulā, obtusā; anfractibus 2, ultimo ventroso; columellā strictiusculā, obliquē recedente; aperturā rotundato-ovali.*

Long. 8, lat.  $6\frac{1}{2}$ , alt. 4 mill.; apert.  $7\frac{1}{2}$  mill. longa.

From the island of Massafuera, Chile (H. Cuming).

4. **SUCCINEA TAHITENSIS**, Pfr. *Succ. testā ovatā, striatulā, tenui, vir nitidā, pellucidā, pallidē succined; spirā brevi, obtusiusculā; anfractibus  $2\frac{1}{2}$  convexis, ultimo ovato; columellā leviter arcuatā, medio obsoletē (interdum distinctē) angulatā; aperturā regulariter ovali; peristomate expansiusculo.*

Long. 12, lat. 7, alt.  $4\frac{1}{2}$  mill.; apert.  $8\frac{2}{3}$  mill. longa.

From Tahiti (H. Cuming).

5. **HELIX LINDONI**, Pfr. *Hel. testā imperforatā, semiglobosā, tenui, irregulariter striatulā, supernē opacā, albida, punctis castaneis conspersā et linea nigricanti-castaneā ad suturam ornatā; anfractibus  $4\frac{1}{2}$  convexiusculis, ultimo basi planulato, pellucido, corneō-virente, anticē breviter deflexo; columellā intrante, declivi, subarcuatā, dilatatā, introrsum acutā; aperturā lunato-ellipticā; peristomate simplece, recto.*

Diam. 16, alt.  $9\frac{1}{2}$  mill.

From the island of Cuba (Lindon).

6. **HELIX PEMPHIGODES**, Pfr. *Hel. testā imperforatā, subglobosā, tenui, membranacea, obliquē plicatulā, diaphanā, lutescenti-corneā; spirā brevi, papillatā; anfractibus 4 vix convexis, ultimo permagno, carinato, juxta suturam inflato, basi convexo, anticē vix descendente; columellā simplice, acutā, subverticali; aperturā amplā, ferē circulari; peristomate simplice, acuto, marginibus conniventibus.*

Diam. 18, alt. 12 mill.

From the island of Cuba (Lindon).

7. **HELIX GRADATA**, Pfr. *Hel. testā imperforatā, globoso-turbanatā, striatulā, tenui, hyalina; spirā turbinatā, ad apicem acutā; anfractibus 6 convexiusculis, gradatis, ultimo medio acutē carinato, basi convexo, sub lente minutissimē concentricē striato; aperturā sub-tetragono-lunari; peristomate simplice, acuto, margine columellari verticaliter descendente.*

Diam. 5, alt. 5 mill.

From the island of Leyte (H. Cuming).

Nearly allied to *H. tongana*, Quoy.

8. **HELIX BARCLAYANA**, Pfr. *Hel. testā umbilicatā, depresso-turbanatā, confertim obliquē costatā, albā, epidermide fusco-olivacea indutā; spirā conoidea, apice obtuso; suturā profundā; anfractibus 5½ convexis, ultimo medio carinato (interdum obsoletē bicarinato), basi convexiusculo; umbilico mediocri, ferē cylindrico; aperturā subverticali, quadrangulari; peristomate simplice, acuto, margine columellari verticaliter descendente, cum basali angulum formante.*

Diam. 18, alt. 12 mill.

From the island of France (Sir D. Barclay).

9. **HELIX ARCUATA**, Pfr. *Hel. testā umbilicatā, orbiculato-convergē, tenui, pellucidā, pallidē cornē, confertim et regulariter arcuatoplicatā; spirā latē conoidea, apice obtusiusculo; anfractibus 6 vix convexiusculis, carinā acutā, serratā marginatis, ultimo circa umbilicum magnum, ferē cylindricum subcompressō; aperturā angulato-lunari, latā; peristomate simplice, acuto, margine columellari brevi, verticali.*

Diam. 5½, alt. 2¾ mill.

From the province of Cagayan, island of Luzon (H. Cuming).

10. **HELIX MIGHELSIANA**, Pfr. *Hel. testā umbilicatā, globosā, solidā, validē et confertim plicatā, spiraliter obsoletē striatā, rufā vel lutescenti-fuscā; spirā conoideo-semiglobosā; anfractibus 5 convergiensculis, ultimo ventroso, anticē vix descendente, circa umbilicum angustum compressō; aperturā subverticali, rotundato-lunari; peristomate recto, intus albo-labiato, margine columellari dilatato-patente.*

Diam. 19, alt. 15 mill.

From Surigao, island of Mindanao (H. Cuming).

11. **HELIX RISSOANA**, Pfr. *Hel. testā perforatā, globosā, tenui,*

*striatd, diaphaná, vix nitidá, rufd; spird conoided, obtusiusculd; anfractibus 6 convexiusculis, ultimo subangulato, medio pallidé cingulato, anticè breviter descendente, basi ventroso; aperturd magnd, semicirculari; peristomate intus rubello-labiato, breviter expanso, margine columellari in laminam brevem, perforationem semioccultantem reflexo.*

Diam. 18, alt. 13 mill.

From Greece (Lieut. Spratt, R.N.).

12. *HELIX DICTYODES*, Pfr. *Hel. testá angustè umbilicatá, depressá, sublenticulari, obliquè plicato-striatd, tenuiusculd, diaphaná, pallidé corned, fusco subtiliter reticulatá et maculis castaneis juxta suturam et carinam ornatá; spird latè conoided; anfractibus 7 vix convexiusculis, ultimo acutè carinato; aperturá subverticali, depressd, lunari, intus margaritaced; peristomate simple, margine columellari breviter dilatato-patente, basali sinuoso, reflexiusculo.*

Diam. 27, alt. 12 mill.

From New Guinea (Ince).

13. *HELIX LIGNARIA*, Pfr. *Hel. testd imperforatd, subgloboso-depressd, solidá, lavigatd, castanéa, fasciis variis epidermidis hydrophanæ, fusco-cinereæ obductd; spird vix elevatá, obtusá; anfractibus 4½ vix convexiusculis, celeriter accrescentibus, ultimo ad peripheriam subangulato; columelld strictiusculd, peroobliqud, latd, planatd, fuscá; aperturd rotundato-lunari, intus albida; peristomate subincrassato, brevissimè reflexo, fusco-marginato.*

Diam. 45, alt. 27 mill.

From Surigao, island of Mindanao (H. Cuming).

14. *HELIX CRASSILABRIS*, Pfr. *Hel. testd imperforatd, depressd, crassd, ponderosá, irregulariter striatd et undique granulatd, albida, lineis spiralibus, undulatis, fuscis, fascidque unid ad peripheriam ornatd; spird vix elevatd, distinctè granulatá, apice nudo, albo; anfractibus 4½ planiusculis, sensim accrescentibus, ultimo minutissimè granulato, medio subcarinato, anticè vix descendente; aperturd obliqud, semielliptica, intus albá; peristomate undique incrassato-reflexo, margine dextro subsinuoso, columellari intus obsoletè plicato.*

Diam. 42, alt. 22 mill.

From the island of Cuba (Lindon).

15. *HELIX SPENGLERIANA*, Pfr. *Hel. testd imperforatd, depressd, solidá, striatd, nitidá, pallidé castaned; spird parum elevatá, obtusd; anfractibus 5½, supremis planis, minutissimè granulatis, 2 ultimis convexis, ultimo medio obtuse carinato, basi convexiusculo; aperturd peroobliqud, lunari, intus fusculá; peristomate latè expanso, breviter reflexo, marginibus callo nitido junctis, basali sinuoso, reflexo, subappresso, columellari perdilatato, adnato, umbilicum prorsus tegente.*

Diam. 49, alt. 26 mill.

From the island of Jamaica (Gosse).

16. **HELIX CODONODES**, Pfr. *Hel. testā umbilicatā, globoso-conoidē, solidā, obliquè striatulā, lineis spiralibus confertis subtilissimè sculptā, nitidā, albā, castaneo-bifasciatā; spirā campanulatā, apice obtusiusculo; anfractibus 5½ vix convexis, ultimo anticè deflexo, basi juxta aperturam gibboso-subconstricto; aperturā obliquā, ferè circulari; peristomate incrassato, reflexo, marginibus approximatīs, callo nitido junctis, columellari dilatato, patente, sinuoso.*

Diam. 20, alt. 17 mill.

From the Philippine Islands (H. Cuming).

The described specimen shows a tooth-like protuberance on the inner side of the columella, which seems not to belong to the essential characters of this species.

17. **BULIMUS CASTUS**, Pfr. *Bul. testā subperforatā, ovato-conicā, tenuiusculā, minutim et obsoletè decussatulā, hyalino-albiddā, basi et prope aperturam erubescens; spirā conicā, acutiusculā; anfractibus 5½ convexiusculis, ultimo spiram paulo superante; columellā strictiusculā, filiformi; aperturā oblongā; peristomate simplice, roseo, marginibus subparallelis, callo tenui junctis, dextro breviter expanso, columellari brevissimè reflexo, perforationem ferè claudente.*

Long. 19, diam. 9 mill.

From Central America? (Latre).

18. **BULIMUS ERUBESCENS**, Pfr. *Bul. testā subperforatā, oblongo-turritā, ləviusculā, lineis spiralibus sub lente insculptā, carneolata, apice rubicundā; spirā turritā, apice acuto; anfractibus 6 planiusculis, ultimo spirā paulo breviore; columellā supernè subtortā, basi paulo recedente; aperturā oblongā, intus nitide albd; peristomate simplice, margine dextro expansiusculo, columellari forniciatim breviter reflexo, subappresso.*

Long. 24, diam. 10 mill.

Locality unknown.

19. **BULIMUS RIMATUS**, Pfr. *Bul. testā profundè rimatā, oblongo-turritā, tenuiusculā, subarcuatim striatulā, pallidè corned; spirā turritā, obtusiusculā; anfractibus 7 ferè planis, ultimo ¾ longitudinis æquante, basi rotundato; columellā intus uniplicatā; aperturā oblongo-ovali; peristomate simplice, marginibus approximatīs, callo junctis, dextro vix expanso, columellari dilatato, patente.*

Long. 33, diam. 11 mill.

Locality unknown.

20. **BULIMUS STUDERI**, Pfr. *Bul. testā perforatā, oblongo-conicā, tenuiusculā, striatulā, lineis spiralibus confertis sub lente decussatā, nitidā, albā, cingulis angustis, roseis 3-4 ornatā; spirā conicā, acutā; anfractibus 6 vix convexiusculis, ultimo ¾ longitudinis subæquante; columellā arcuatā; aperturā ovali-elliptica, intus colore; peristomate simplice, marginibus subconniventibus, dextro breviter expanso, columellari forniciatim reflexo, roseo.*

Long. 25, diam. 10 mill.

From Central America? (H. Cuming.)

21. **BULIMUS MORICANDI**, Pfr. *Bul. testā perforatd, ovato-conicā, tenui, lineis spiralibus subconfertis insculptd, subdiaphand, citrind; spirā conicā, acutiusculā; suturā pallidē submarginatd; anfractibus 6 vix convexis, ultimo spiram æquante; columella strictid; aperturd suboblongā, truncato-ovali, intus concolore; peristomate simplice, breviter expanso, margine columellari supernè breviter patentī-reflexo.*

Long. 24, diam. 12 mill.

From Mount Coban, Central America (Latre).

22. **BULIMUS EHRENBERGI**, Pfr. *Bul. testā profundē rimatā, oblongā, solidulā, obliquē striatuld, albida; spirā oblongā, apice attenuato, obtusiusculo; anfractibus  $7\frac{1}{2}$  vix convexiusculis, ultimo  $\frac{2}{5}$  longitudinis pauld superante; aperturā angulato-ovali; peristomate incrassato, breviter reflexo, marginibus callo crasso, prope insertionem labri tuberculifero junctis, columellari dilatato, crasso, patente.*

Long. 24, diam. 10 mill.

From Cerigotto, Greece (Lieut. Spratt, R.N.).

23. **BULIMUS ROSSMASSLERI**, Pfr. *Bul. testā profundē rimatā, oblongā, solidulā, confertim rugoso-plicatā, supernē fusco-corneā, basi sordidē albida; spirā oblongo-conicā, apice obtuso; anfractibus 8 vix convexiusculis, ultimo basi rotundato,  $\frac{1}{3}$  longitudinis æquante; columellā brevi, strictiusculā; aperturd truncato-ovali, intus albā; peristomate albo-labiato, breviter expanso, marginibus callo tenui, juxta insertionem labri dentifero junctis, columellari dilatato, patente.*

Long. 19, diam. 7 mill.

Locality unknown.

24. **BULIMUS DRAPARNAUDI**, Pfr. *Bul. testā subobtectē perforatā, oblongo-subfusiformi, striatuld, opacā, nitidā, albā, cærulescenti-nebulosā, strigis nigro-castaneis et brunneis, interdum maculosē interruptis, ornatā; spirā turrito-conicā, ad apicem acutā; anfractibus 7 convexiusculis, ultimo  $\frac{3}{8}$  longitudinis subæquante; columella rectā; aperturā oblongā; peristomate simplice, acuto, margine columellari dilatato, membranacea, angulatim reflexo, appresso.*

Long. 28, diam. 11 mill.

*β. Minor, interstitiis strigarum castaneo-litturatis.*

From Chilon, Bolivia (Bridges).

25. **BULIMUS ZIEGLERI**, Pfr. *Bul. testā subperforatd, ovato-conicd, tenui, confertim striatuld, lineis spiralibus sub lente obsoletē decussatd, albida; spirā conicā, acutiusculā; anfractibus 6 vix convexiusculis, ultimo medio subangulato, spirā pauld breviore; columella pauld recedente; aperturd ovali; peristomate simplice, margine columellari breviter reflexo, subappresso.*

Long. 21, diam. 10 mill.

*β. T. pellucidā, lutescente, fasciis castancis, supremis maculosē interruptis, cinctā.*

Locality unknown.

26. **BULIMUS SAYI**, Pfr. *Bul. testā subperforatā, ovato-oblonga, solidiuscula, confertim rugoso-plicata, nitida, albā, strigis pellucidis, fuscis ornata; spirā conicā, obtusiuscula; anfractibus 6 vix convexis, ultimo spiram subēquante, basi attenuato, circa perforationem obsoletam fusco-areolato; columella leviter arcuatā; aperturā elliptico-oblonga, intus fusco-carneū; peristomate simple, margine columellari breviter reflexo, subappresso.*

Long. 20, diam. 9 mill.

Locality unknown.

27. **BULIMUS CONIFORMIS**, Pfr. *Bul. testā subperforatā, ovato-conicā, tenui, irregulariter striatā, fuscescenti-albida, strigis obliquis, fuscis signata; spirā conicā, acutiuscula; anfractibus 5 planiusculis, ultimo spiram paulo superante, medio angulato, basi subcompresso; columella leviter arcuatā; aperturā ovali, utrinque angustatā; peristomate simple, recto, margine columellari supernè dilatato, breviter reflexo.*

Long. 12, diam. 6½ mill.

From Merida, Andes of Bolivia (T. Bridges).

28. **BULIMUS SOWERBYI**, Pfr. *Bul. testā perforatā, ovato-conicā, tenui, sublævigata, albida, strigis obliquis, castaneis, maculas albas pyramidales et rhomboidales formantibus ornata; spirā conicā, acutā; anfractibus 6½ vix convexiusculis, ultimo spiram aequante, medio pallide, juxta basin attenuatam castaneo-unifasciato; columella paulo recedente; aperturā oblongo-ovali; peristomate simple, recto, margine columellari angulatim latè reflexo, plano.*

Long. 22, diam. 10 mill.

From the Columbian Andes (Lindon).

29. **BULIMUS PORPHYRIUS**, Pfr. *Bul. testā perforatā, oblongo-attenuata, solidiuscula, confertim et ruditer corrugata, castanea, strigis albis irregulariter marmorata; spirā conicā, ad apicem obtusa; suturā submarginata, irregulariter crenata; anfractibus 7 planiusculis, summis subtiliter granulatis, ultimo spirā paulo breviore; columella subrecta; aperturā angusta, oblonga; peristomate simple, acuto, margine columellari dilatato, reflexo, carneolivido, perforationem ferè occultante.*

Long. 51, diam. 20 mill.

From Bolivia (T. Bridges).

30. **BULIMUS VOITHIANUS**, Pfr. *Bul. testā perforatā, subfusiformis oblonga, solidula, rugis longitudinalibus et lineis concentricis impressis ruditer granulata, sordidè albā; spirā conicā, ad apicem acutiuscula; anfractibus 6–7 vix convexiusculis, ultimo spirā paulo breviore; columella subverticali, nigro-castanea; aperturā angusta, oblonga, intus castanea; peristomate simple, recto, marginibus callo fusco junctis, columellari dilatato, fornicateum reflexo, perforationem profundam non tegente.*

Long. 19, diam. 7½ mill.

From Chile (T. Bridges).

31. **BULIMUS CASTRENSIS**, Pfr. *Bul. testā angustē umbilicatd, oblongo-conicd, lēvissimē striatuld, opacd, albidā, strigis spadiceis denticulatis et maculis albis pyramidalibus ornatd; spirā conicd, acutiusculd; anfractibus 7 vix convexiusculis, ultimo ventrosiore, infra medium lineis nonnullis spadiceis cincto,  $\frac{2}{5}$  longitudinis subæquante; columellā strictiusculd; aperturd oblongā; peristomate simplice, recto, margine dextro supernē arcuato, columellari dilatato, patente.*

Long. 19, diam. 9 mill.

Locality unknown.

32. **BULIMUS ANDICOLA**, Pfr. *Bul. testd perforatd, turrito-conicd, solidā, lineis concentricis, confertis sub lente sculptā, opacd, nitidā, albā, strigis fuscis, linearibus irregulariter ornatā; spirā elongatd, acutiusculd; anfractibus 7 convexiusculis, ultimo  $\frac{3}{7}$  longitudinis subæquante, basi rotundato; columellā deorsum aliquantulum rececente; aperturā ovali-oblongā; peristomate simplice, acuto, margine columellari supernē fornicatim reflexo, perforationem angustum formante.*

Long. 24, diam. 11 mill.

From the Columbian Andes (Lindon).

33. **PUPA ELEGANTULA**, Pfr. *Pup. testā breviter rimatā, subcylindraceā, apice obtuso, lēvigato, nitido, hyalino; anfractibus 7 planiusculis, ultimo præcedente pauld angustiore, extus medio sulcato, intus lamellis 2 validis, suturæ parallelis, plicuque profundd columellæ paralleld munito; aperturā subsemicirculari, lamellā parietis aperturalis intrante juxta insertionem labri coarctatā; peristomate expansiusculo, margine dextro flexuoso, medio subincrassato.*

Long. 6 $\frac{2}{3}$ , diam. 3 mill.

Locality unknown.

34. **ACHATINA LAMARCKIANA**, Pfr. *Ach. testd ovato-conicd, solidā, ponderosā, ruditer plicatā, in fundo albido strigis fulminatis nigritibus et castaneis, maculisque rufis variegatd; spirā conicd, pallida, apice obtusiusculo; anfractibus 8 convexiusculis, supremis lineis spiralibus obsoletē decussatis, ultimo ventroso, spiram superrante; columellā arcuatā, purpureo-callosā, supra basin aperturæ obliquè et leviter truncatd; aperturā ovali, intus margaritacea, cœrulescente, saturatius marmorata; peristomate fusco-limbato, marginibus callo purpureo junctis.*

Long. 103, diam. 52 mill.

From the interior of the island of Madagascar.

35. **ACHATINA RANGIANA**, Pfr. *Ach. testā elongatā, turritā, solidā, ponderosā, lēvissimē arcuatim substriatā, lineis spiralibus distan- tibus notatā, stramineā, apice albo, obtusiusculo; suturā lēvissimā; anfractibus 11 planulatis, ultimo  $\frac{1}{4}$  longitudinis pauld superante, basi rotundato; columellā rectā, callosā, ad basin aperturæ breviter et obliquè truncatā; aperturā subsemiovali, intus margaritacea; peristomate simplice, acute.*

Long. 39, diam. 11 mill.  
From Mexico (Lindon).

36. *ACHATINA BULIMOIDES*, Pfr. *Ach. testa ovato-conicā, tenui, striatā, epidermide corneo-luted, pellucida indutā; spirā conicā, acutā; anfractibus 5½ vix convexis, ultimo ventrosiore, spiram æquante; columellā supernè tortā, filari, supra basin aperturæ obsoletissimè truncatā, callo tenui ventrem anfractū penultiimi vestiente munitā; aperturā latè semiovali; peristomate simple, tenui.*

Long. 11, diam. 6 mill.

From the island of Juan Fernandez (H. Cuming).

37. *ACHATINA (GLANDINA) LINDONI*, Pfr. *Ach. testa oblonga, utrinque attenuatā, solidulā, lævigatā, nitida, pallidè fulva, lineis incrementi arcuatis, vix prominētibus, saturationibus notata; spirā conicā, acutiusculā; suturā submarginatā; anfractibus 8 planiusculis, 2 ultimis obliquè descendantibus, ultimo spiram æquante, supra columellam intus gibboso; columella brevi, ad basin aperturæ obliquè truncatā; aperturā angustissimā, basi subcanaliculatā; peristomate simple, marginibus callo junctis, dextro antrorum arcuato-dilatato.*

Long. 21, diam. 6 mill.

From the island of Cuba (Lindon).

38. *CYLINDRELLA SOWERBYANA*, Pfr. *Cyl. testa truncatā, cylindraceo-subulatā, solidiusculā, obliquè subarcuatim costulato-striatā, opacā, cinnamomeo et albo radiatā; anfractibus (spec. trunc.) 16 angustis, convexiusculis, ultimo basi subcarinato (carina parum prominente, ferè rectangula), anticè vix protracto, subtilius striato; aperturā subcirculari; peristomate undique libero, tenui, breviter expanso, margine supero sursum dilatato.*

Long. 35, diam. 8 mill.

From the island of Cuba (Lindon).