

PROCEEDINGS

OF THE

ZOOLOGICAL SOCIETY

OF LONDON.



PART XXIV.

1856.

PRINTED FOR THE SOCIETY ;
SOLD AT THEIR HOUSE IN HANOVER-SQUARE,
AND BY MESSRS. LONGMAN, BROWN, GREEN AND LONGMANS,
PATERNOSTER-ROW.

4. DESCRIPTIONS OF TWENTY-SEVEN NEW SPECIES OF LAND-SHELLS, COLLECTED BY M. SALLÉ IN THE STATE OF VERA CRUZ, MEXICO. BY DR. L. PFEIFFER.

(Mollusca, Pl. XXXV.)

1. *HELIX CORDOVANA*, Pfr. *H. testa umbilicata, deppressa, tenui, undique subgranulata et breviter pilosa, cornea, superne fasciis 2 angustis, rufis cincta; spira plana; sutura profunda, canaliculata; anfr. 4½ turgidis, sensim accrescentibus, ultimo supra peripheriam obsolete sulcato, antice descendente, subitus rotundato; umbilico ¼ diametri subæquante; apertura perobliqua, lunato-circulari; perist. tenui, marginibus conniventibus, supero expanso, basali reflexiusculo.*

Diam. maj. $12\frac{1}{3}$, min. $10\frac{1}{3}$, alt. 5 mill.

Hab. Cordova.

2. *HELIX VERACRUZENSIS*, Pfr. *H. testa umbilicata, deppressa, tenerrima, striatula, pellucida, nitidissima, pallide rubello-cornea; spira parum elevata, vertice subtili; sutura subcrenulato-marginata; anfr. 5 rix convexiusculis, ultimo lato, non descendente, depresso-rotundato; umbilico perrivo, $\frac{1}{6}$ diametri vix æquante; apertura obliqua, lunato-ovali; perist. recto, acuto, marginibus subconniventibus, columellari arcuato-declivi, vix reflexiusculo:*

Diam. maj. $12\frac{1}{2}$, min. $10\frac{1}{2}$, alt. 5 mill.

Hab. Cordova.

3. *BULIMUS SULPHUREUS*, Pfr. (Pl. XXXV. f. 11.) *B. testa anguste umbilicata, ovato-conica, tenuiuscula, striatula, striis spiralibus confertissimis decussata, nitida, pallide sulphurea; spira conica, apice acuta, concolore; anfr. 6½ convexiusculis, ultimo spira breviore, basi rotundato; columella subprecedente; apertura obliqua, ovali; perist. simplice, tenui, margine dextro breviter expanso, superne sinuato, columellari triangulatim dilatato, reflexo.*

Long. 29, diam. 12 mill.

Hab. Cordova.

4. *BULIMUS CORIACEUS*, Pfr. *B. testa anguste perforata, ovato-conica, solidula, sub lente obsolete granulata, fusco-cornea, haud nitente; anfr. 6 convexiusculis, summis nigro-fasciis, sequentibus castaneo-fasciatibus, ultimo spira breviore, obsolete angulato, basi subattenuato; columella arcuata; apertura parum obliqua, elliptico-ovali; perist. simplice, recto, margine columellari albido, nitido, fornicatim reflexo.*

Long. 18, diam. 9 mill.

Hab. Cordova.

5. *BULIMUS MARTENSI*, Pfr. *B. testa subperforata, turrito-oblonga, tenui, striatula et distanter chordato-costata, diaphana, albido-hyalina; spira turrita, obtusula; anfr. 6 convexis, ulti-*





mo $\frac{2}{5}$ longitudinis vix æquante, rotundato; columella medio subdentato-plicata; apertura parum obliqua, elliptico-ovali; perist. simplice, recto, margine columellari late reflexo, sublibero.
Long. $9\frac{1}{3}$, diam. $4\frac{1}{2}$ mill.
Hab. Cordova.

6. **BULIMUS COSTATO-STRIATUS**, Pfr. *B. testa imperforata, turrita, tenui, conferte striata et costulis irregularibus munita, diaphana, cereo-hyalina; spira regulariter attenuata, acutiuscula; anfr. 7 convexiusculis, ultimo $\frac{2}{7}$ longitudinis formante, basi rotundato; columella substriata, filari; apertura vix obliqua, oblonga; perist. simplice, recto, marginibus subparallelis, columellari simplice.*

Long. $7\frac{1}{2}$, diam. $2\frac{1}{2}$ mill.
Hab. Cordova.

7. **BULIMUS DROUËTI**, Pfr. (Pl. XXXV. f. 12.) *B. testa subobtecta perforata, ovato-conica, tenui, ruguloso-striata (striis spiralibus obsoletissime decussata), pallide straminea, strigis et fasciis latis spadiceis interruptis ornata; spira conica, acutiuscula; anfr. 6 convexiusculis, ultimo spiram vix superante, antice subvaricosa; columella substricta, compressa; apertura obliqua, ovali; perist. tenui, expansiusculo, margine columellari dilatato, abrupte reflexo.*

Long. 24, diam. $10\frac{1}{2}$ mill.
Hab. Cordova.

8. **BULIMUS AURIFLUUS**, Pfr. (Pl. XXXV. f. 10.) *B. testa perforata, ovato-conica tenui, sub lente minutissime decussata, nitida, albida, strigis angustis flexuosis fulvo-aureis ornata; spira convexiusculo-conica, acuta; anfr. $5\frac{1}{2}$ convexiusculis, ultimo spiram paulo superante, juxta perforationem angustum subattenuato, unicolor lutescente; columella intrante, leviter arcuata; apertura vix obliqua, oblonga; perist. tenui, breviter expanso, margine columellari compresso, superne dilatato, reflexo.*

Long. 22, diam. 10 mill.
Hab. Cordova.

9. **SIMPULOPSIS SALLEANA**, Pfr. (Pl. XXXV. f. 15, 16.) *S. testa subsemiovata, solidula, striatula, lineisque spiralibus impressis obsoletis notata, nitida, corneo-straminea; spira parvula, obtusa; anfr. vix $2\frac{1}{2}$, ultimo magno, inflato; columella arcuata, subacuta; apertura pero obliqua, lunato-rotundata, intus margaritacea; perist. simplice, marginibus callo tenuissimo junctis, dextro expansiusculo, antrorsum dilatato.*

Diam. maj. 15, min. 12, alt. $7\frac{1}{2}$ mill.
Hab. Cordova.

10. **SIMPULOPSIS CORDOVANA**, Pfr. *S. testa subsemiglobosa, tenuissima, levissime striatula, pellucida, nitidissima, virenti-*

cornea; spira minuta, vix prominula; sutura canaliculata; anfr. 2½, ultimo inflato; columella tenui, papyraceo-marginata; apertura pero obliqua, fere circulari; perist. simplex, recto, margine dextro superne antrorsum dilatato.

Diam. maj. 15, min. 11½, alt. 7 mill.

Hab. Cordova.

11. **SPIRAXIS SHUTTLEWORTHI**, Pfr. (Pl. XXXV. f. 8.) Sp. testa oblongo-glandiformi, tenuiuscula, laevigata, lineis impressis irregularibus notata, lucida, corneo-flavescens, strigis variciformibus virenti-fusca notata; spira conica, obtusula; sutura levi, sublacerata; anfr. 7, superis vix convexiusculis, ultimo antice descendente, $\frac{2}{3}$ longitudinis fere aequante, basi subattenuata; lamina columellari leviter torta, subincurvata, basi vix truncatula; apertura anguste semiovali, superne longe acuminata, intus margaritacea; perist. simplex, marginibus callo tenui junctis, dextro medio dilatato.

Long. 33, diam. 13½ mill.

Hab. Cordova.

12. **SPIRAXIS TURGIDULA**, Pfr. (Pl. XXXV. f. 9.) Sp. testa ovato-conica, tenuiuscula, striatula, nitida, pallide flavescens, strigis variciformibus, subimpressis, pellucidis notata; spira conica, obtusiuscula; sutura crenulata; anfr. 8 infra suturam turgidulis, ultimo $\frac{4}{7}$ longitudinis subaequante; lamina columellari crassa, alba, leviter torta, basi truncata; apertura subverticale, sinuoso-semiovali, intus albido-margaritacea; perist. simplex, margine dextro medio antrorsum dilatato.

Long. 31, diam. 12½ mill.

Hab. Cordova.

13. **SPIRAXIS AURICULACEA**, Pfr. Sp. testa fusiformi-oblonga, tenera, laevigata, pellucida, nitida, rubello-cornea; spira elongato-conica, obtusula; sutura marginata; anfr. 7 convexiusculis, ultimo spira vix longiore, infra medium dilatato; lamina columellari parum torta, alba, filari, minime truncata; apertura sinuato-semiovali, intus leviter margaritacea; perist. simplex, margine dextro medio fere angulatum producto.

Long. 16, diam. 6 mill.

Hab. Cordova.

14. **ACHATINA (VARICELLA) ORIZABÆ**, Pfr. (Pl. XXXV. f. 6.) A. testa oblongo-conica, solidula, longitudinaliter striata, nitida, olivaceo-fusca, varicibus subprominulis pallidis irregulariter munita; spira elongato-conica, apice obtusula; sutura levissime marginata; anfr. 7½ convexiusculis, ultimo spira breviore, deorsum laeviore; columella subcallosa, arcuata, oblique truncata; apertura verticali, sinuato-ovali; perist. recto, margine dextro obtuso, vix flexuoso, pallide limbato.

Long. 42, diam. 18 mill.

Hab. In Vulcano Orizaba.

15. ACHATINA (VARICELLA) SPECIOSA, Pfr. (Pl. XXXV. f. 7.)

A. testa conico-ovata, solidula, longitudinaliter conferte plicata, nitida, carnea, varicibus sulciformibus, albidis, subflexuosis, irregulariter distantibus munita; spira conica, apice acutiuscula; sutura eleganter nodulato-crenata; anfr. 8 vix convexiusculis, ultimo spiram paulo superante, prope suturam turgidulo, deorsum sublævigato; columella arcuata, late truncata; apertura verticali, sinuato-semiovali; perist. recto, margine dextro flexuoso.

Long. 29, diam. $13\frac{1}{2}$ mill.

Hab. Cordova.

16. ACHATINA (VARICELLA) CORDOVANA, Pfr. *A. tes taturito-oblonga, solidula, longitudinaliter plicata, nitida, alabastrina; spira elongato-conica, apice obtusa; sutura marginata, conferte nodulata; anfr. $7\frac{1}{2}$ parum convexis, ultimo spira paulo breviore, varicibus nonnullis impressis, obsoletis munito, basi vix attenuato; columella substricta, transverse truncata; apertura subverticali, sinuato-semiovali; perist. simplice, margine dextro medio fere angulatim producto, subinflexo.*

Long. 20, diam. 6 mill.

Hab. Cordova.

17. ACHATINA MARGARITACEA, Pfr. *A. testa oblonga, tenui, sublævigata, sub lente levissime et irregulariter plicatula, pellucida, nitida, lutescenti-hyalina; spira brevi, conica, acutiuscula; sutura subcanaliculata, marginata; anfr. $5\frac{1}{2}$ convexiusculis, ultimo $\frac{2}{3}$ longitudinis superante, basi vix attenuato; columella leviter arcuata, abrupte truncata; apertura verticali, acuminate-semiovali; perist. simplice, margine dextro leviter antrorsum arcuato.*

Long. $9\frac{1}{2}$, diam. $3\frac{1}{3}$ mill.

Hab. Cordova.

18. ACHATINA AMBIGUA, Pfr. *A. testa imperforata, ovato-conica, solida, lævigata, opaca, albida; spira conica, obtusula; anfr. 7 convexis, ultimo spira paulo breviore, basi rotundato; columella verticali, subintorta, subtruncata; apertura verticali, trapezio-ovali; perist. recto, margine dextro subrepando, columellæ parallelo.*

Long. 22, diam. 10 mill.

Hab. Cordova.

19. CYLINDRELLA BOUCARDI (Sallé, MSS.), Pfr. (Pl. XXXV. f. 1.) *C. testa arcuato-rimata, clavato-cylindracea, truncata, tenuiuscula, costulis capillaribus, confertis, subarcuatis munita, in interstitiis sub lente transverso-striata, corneo-fulva; sutura levi, vix marginata; anfr. superst. 9-11 vix convexis, ultimo antice soluto, oblique descendente, dorso angulato, infra medium obtuse carinato; apertura angulato-oblonga, plica*

No. CCCXI.—PROCEEDINGS OF THE ZOOLOGICAL SOCIETY.

valida, torta columellæ coarctata; perist. albo, breviter expanso.

Long. 52–56, diam. 13 mill.

Hab. Cordova.

20. **CYLINDRELLA APISTOMA**, Pfr. (Pl. XXXV. f. 4, 5.) *C. testa subrimata, subulata, subareuatim striatula, diaphana, albido-cornea; spira regulariter attenuata, apice integra, acutiuscula; anfr. 22–24 convexiusculis, ultimo breviter protracto, dorso angulato, antice distinctius striato; apertura subverticale, oblique piriformi; perist. albo, undique reflexiusculo, margine dextro superne subsinuoso.*

Long. 17, diam. $2\frac{1}{3}$ mill.

Hab. Cordova.

21. **CYLINDRELLA POLYGYRA**, Pfr. (Pl. XXXV. f. 2, 3.) *C. testa profunde rimata, subulata, gracili, costulis filariibus, confertis, leviter arcuatis sculpta, opaca, cornea; spira regulariter attenuata, apice integra, acutiuscula; anfr. 24–27 convexis, ultimo breviter soluto, dorso et basi subcompresso; apertura vix obliqua, subcirculari, in fundo subtriangulari; perist. undique expanso et reflexiusculo.*

Long. $17\frac{1}{2}$ – $21\frac{1}{2}$, diam. $2\frac{1}{2}$ mill.

Hab. Cordova.

22. **PROSERPINA (CERES) SALLEANA**, Cuming. (Pl. XXXV. f. 21, 22.) *Pr. testa imperforata, conoidea, solidula, superne striis incrementi et granulis minutis exasperata, lutea vel rosea, epidermide opaca, albida, decidua, partim obducta; spira conexo-conoidea, mucronata; anfr. 8 vix convexiusculis, ultimo non descendente, medio compresse et acute carinato, basi convexo, levigato, callo nitido, luteo magis minusve abducto; apertura perobliqua, subtriangulari, lamellis 6 coarctata; parietalibus 2, columellari 1 subtorta, 3 in pariete basali, mediana maxima; perist. luteo, subincrassato, obtuso.*

Diam. maj. 23, min. 21, alt. fere 12 mill.

Hab. Cordova.

23. **HELICINA HELOISE**, Sallé, MSS. (Pl. XXXV. f. 17.) *H. testa turbinato-globosa, solidula, levissime striatula, striis spiraliibus sub lente obsolete decussata, nitida, subunicolore lutea vel cingulo purpurascente, sursum diluto ornata; spira convexo-conoidea, acuminata; sutura pallida; anfr. 6 convexis, ultimo peripheria vix subangulato, basi planiusculo, callo nitido, circumscripto munito; columella arcuata, filari; apertura diagonali, subsemicirculari; perist. tenuiusculo, expanso, margine dextro leviter flexuoso.*

Operc.?—Diam. maj. $9\frac{2}{3}$ –11, min. $8\frac{1}{2}$ – $9\frac{1}{2}$, alt. 7–8 mill.

Hab. Cordova.

24. **HELICINA NOTATA**, Sallé, MSS. (Pl. XXXV. f. 18, 19, 20.)

H. testa globoso-turbinata, solidula, striata et liris subdistantibus, levibus circumdata, carneus vel stramineus; spira convexiusculo-conica, acuta; anfr. $5\frac{1}{2}$ vix convexiusculis, ultimo spiram subaequante, convexiore; columella brevi, basi tuberculata, callum crassum, circumscripsum retrorsum emitente; apertura fere diagonali, subsemicirculari; perist. calloso, incrassato, albo, angulatum patente. — Operc. tenue, margine externo purpureo.

Diam. maj. $8\frac{1}{3}$, min. 7, alt. 7 mill.

Hab. Cordova.

25. **HELICINA CORDILLERÆ**, Sallé, MSS. *H. testa depresso-globoso-conica, solida, striatula, opaca, albida, fascia 1 rubra supra peripheriam, nonnullisque pallidioribus, obsoletis cincta; spira conoidea, sursum fusco-carnea, apice acuta; anfr. $5\frac{1}{2}$ vix convexiusculis, ultimo depresso rotundato; columella extus vix impressa, callosa, basi in nodulum terminata, callum emitente crassiusculum, circumscripsum; apertura diagonali, trianguli-semiorvali; perist. calloso-incrassato, angulatum patente, margine basali in tuberculum columellæ transeunte. — Operc. corneum, castaneum.*

Diam. maj. $12\frac{1}{3}$, min. $10\frac{2}{3}$, alt. 8 mill.

β. Paulo major, unicolor fusco-carnea, impressione columellari distinctiore.

Hab. In monte Orizaba, 12,000' supra Oceanum.

26. **CYCLOSTOMA (CYCLOPHORUS) BOUARDI**, Sallé, MSS. (Pl. XXXV. f. 25.)

C. testa latiuscula umbilicata, conoideo-depressa, solida, impressionibus malleatis undique tuberculato-rugosa et liris obsoletis distantibus munita, epidermide fulva, saturatius fasciata vestita; spira breviter conoidea, obtusa; anfr. 5 modice convexis, celeriter accrescentibus, ultimo rotundato, antice interdum breviter soluto; apertura parum obliqua, subangulato-ovali, intus margaritacea; perist. simplice, recto, continuo, superne obsolete angulato, margine columellari leviter arcuato. — Operc. tenuissimum, fulrum, planum.

Diam. maj. 36, min. 30, alt. 18-19 mill.

Hab. Cordova.

27. **CYCLOSTOMA (CHONDROPOMA) CORDOVANUM**, Pfr. *Ch.*

testa rimato-perforata, turrita, integra, tenui, longitudinaliter confertim plicata, haud nitente, pallide fulva, fasciis interrupitis castaneis ornata; spira regulariter turrita, sursum plerunque violacea, apice submucillari nitida, cornea; anfr. $7-7\frac{1}{2}$ perconverxis, ultimo non soluto; apertura verticali, oralis; perist. duplex; interno albo, breviter porrecto, externo subdilatato,

horizontaliter patente, concentrice striato, castaneo maculato, superne producto, ad anfr. penultimum subexciso.—Operc. albidum.

Long. 13–15½, diam. 6–7½ mill.

Hab. Cordova.

DESCRIPTION OF PLATE XXXV.

- | | | |
|------|-------------|---|
| Fig. | 1. | Cylindrella Boucardi, <i>Pfr.</i> |
| | 2, 3. | — polygyra, <i>Pfr.</i> |
| | 4, 5. | — apistoma, <i>Pfr.</i> |
| | 6. | Achatina (Varicella) Orizabæ, <i>Pfr.</i> |
| | 7. | (Varicella) speciosa, <i>Pfr.</i> |
| | 8. | Spiraxis Shuttleworthi, <i>Pfr.</i> |
| | 9. | — turgidula, <i>Pfr.</i> |
| | 10. | Bulimus aurifluus, <i>Pfr.</i> |
| | 11. | — sulphureus, <i>Pfr.</i> |
| | 12. | — Drouëti, <i>Pfr.</i> |
| | 13, 14. | Helix caduca and var., <i>Pfr.</i> |
| | 15, 16. | Simpulopsis Salleana, <i>Pfr.</i> |
| | 17. | Helicina Heloisæ, <i>Sallé.</i> |
| | 18, 19, 20. | — notata and var., <i>Pfr.</i> |
| | 21, 22. | Proserpina (Ceres) Salleana, <i>Cuming.</i> |
| | 23, 24. | — eolina, <i>Ductos</i> (for comparison). |
| | 25. | Cyclophorus Boucardi, <i>Sallé.</i> |

5. DESCRIPTIONS OF FIFTY-EIGHT NEW SPECIES OF HELICEA FROM THE COLLECTION OF H. CUMING, ESQ. BY DR. L. PFEIFFER.

1. VITRINA FLEMINGI, *Pfr.* *V. testa subdepressa, peripheria auriformis; solidula, superne plicato-striata striisque spiralibus sub lente notata, cæneo-micante, olivaceo-fulva; spira parum elata; sutura anguste albumarginata; anfr. 4½ convexiusculis, ultimo magno, infra medium obsoletissime angulato, basi levigato, nitidiore; apertura diagonali, lunato-ovali, intus margaritacea; perist. simplice, margine dextro subrepando, columellari arcuato, superne triangulatim reflexo, adnato.*

Diam. maj. 33, min. 24, alt. 17–18 mill.

Hab. Scinde, India (*Dr. Alex. Fleming*).

2. VITRINA PLANTI, *Pfr.* *V. testa deppressa, ambitu subauriformi, tenuissima, levigata, nitidissima, pellucida, albido-hyalina; spira parvula, plana; sutura vix impressa, submarginata; anfr. vix 3 planiusculis, ultimo magno, superne vix convexiore, ad suturam striatulo, subtus inflato, membranaceo-marginato; apertura magna, diagonali, lunato-ovali; perist. simplice, margine dextro antrorsum arcuato, columellari tenuissima.*

Diam. maj. 12, min. 8, alt. 6 mill.

Hab. Natal (*Mr. Plant*).

3. VITRINA BORNEENSIS, *Pfr.* *V. testa depresso-globosa, tenuissima, striatula, striis spiralibus nonnullis impressis notata, pellu-*

cida, pallide cornea; spira brevissime conoidea, albida; sutura submarginata; anfr. 4 celeriter accrescentibus, superis vix convexis, ultimo subdepresso-rotundato, basi oblique ruguloso; apertura obliqua, lunato-rotundata; perist. simplice, levissime inflexo, margine columellari brevi, strictiusculo, filari.

Diam. maj. 15, min. $11\frac{2}{3}$, alt. 9 mill.

Hab. Borneo.

4. **VITRINA CELEBENSIS**, Pfr. *V. testa helicoidea, solidula, striatula, nitida, coffeacea; spira breviter conoidea, obtusula; sutura pallide marginata; anfr. 4 convexiusculis, sensim accrescentibus, ultimo basi parum convexo, pallido; apertura diagonali, lunato-rotundata; perist. simplice, recto, marginibus subconvergentibus, columellari arcuato, subcalloso.*

Diam. maj. $11\frac{1}{2}$, min. 9, alt. 7 mill.

Hab. Celebes (Mrs. Ida Pfeiffer).

5. **VITRINA IDÆ**, Pfr. *V. testa depresso-globosa, ambitu subovali, tenui, arcuato-striatula, nitida, pellucida, lutescenti-cornea; spira vix elevata; sutura levi; anfr. 4 convexiusculis, ultimo magno, inflato, basi non impresso; apertura diagonali, rotundato-lunari; perist. simplice, marginibus conniventibus, regulariter arcuatis, columellari brevissime recedente, subcalloso.*

Diam. maj. 14, min. 11, alt. $7\frac{2}{3}$ mill.

Hab. Celebes (Mrs. Ida Pfeiffer).

6. **VITRINA ? COMORENSIS**, Pfr. *V. testa subglobosa, tenui, rugoso-striata, vix nitidula, diaphana, virenti-cornea, punctis et strigis luteis variegata; spira convessa, obtusa; anfr. 3 convexis, rapide accrescentibus, ultimo inflato; apertura fere diagonali, lunato-ovali, intus margaritacea, albo variegata; perist. simplice, recto, marginibus convergentibus, regulariter arcuatis, columellari filari.*

Diam. maj. $21\frac{1}{2}$, min. $16\frac{1}{2}$, alt. 13–14 mill.

Hab. Mayotte, Comoro Islands (Mr. Clouë).

Shape very similar to that of *Helix aperta*, Born.

7. **VITRINA DARNAUDI**, Pfr. *V. testa semiovata, pertenui, sublente irregulariter striatula punctisque impressis notata, oleosomicante, pallide virenti-cornea; spira convessa, obtusa; sutura anguste marginata; anfr. vix 3 convexis, rapide accrescentibus, ultimo magno; apertura subdiagonali, sublunato-ovali, intus leviter margaritacea; perist. simplice, brevissime inflexo, marginibus approximatis, dextro antrorum dilatato, columellari arcuato, anguste membranaceo-marginato.*

Diam. maj. 16, min. 12, alt. $7\frac{1}{2}$ mill.

Hab. Sennaar, Interior of Africa (Mr. Darnaud).

8. **VITRINA SENNAARIENSIS**, Pfr. *V. testa depressa, tenui, oblique rugosa, nitidula, pellucida, corneo-virente, plerumque luteo*

nigro obducta ; spira subturbinata, mucronata ; anfr. 3 convexis, sensim accrescentibus, ultimo depresso, peripheria obsolete ungu-lato ; apertura perobliqua, lunato-circulari ; perist. tenui, subin-flexo, marginibus convergentibus, columellari vix dilatato.

Diam. maj. $6\frac{2}{3}$, min. $5\frac{1}{3}$, alt. 3 mill.

Hab. Sennaar, Interior of Africa (Mr. Darnaud).

9. **VITRINA LEUCOSPIRA**, Pfr. *V. testa depresse subglobosa, am-bitu ovali, tenuissima, striatula, pellucida, nitida, lutescenti-hy-a-lina ; spira parum convexa, alba ; sutura vix submarginata ; anfr. 4 convexiusculis, penultimo sensim pellucido-radiato, ultimo ro-tunduto ; apertura obliqua, lunato-subcirculari ; perist. tenui, marginibus conniventibus, subinflexis, dextro antrorum valde ar-cuato, columellari substricto, brevi, filari.*

Diam. maj. $13\frac{1}{2}$, min. $11\frac{1}{2}$, alt. $8\frac{1}{2}$ mill.

Hab. Australia.

10. **SUCCINEA PLANTI**, Pfr. *S. testa conico-ovata, tenuiuscula, rugulosa, diaphana, sordide cornea ; spira brevi, subpapillata ; anfr. vix $2\frac{1}{2}$, penultimo conveexo, ultimo $\frac{2}{3}$ longitudinis superante, medio turgido ; columella substricte recedente, superne levissime calloso-plicata ; apertura fere diagonali, ubique incumbente, acu-minato-ovali ; perist. simplice, margine dextro regulariter arcuato.*

Long. 5, diam. $3\frac{1}{2}$, alt. $2\frac{2}{3}$ mill.

Hab. Cape Natal (Mr. Plant).

11. **SUCCINEA ASPERULA**, Pfr. *S. testa ovato-conica, tenuiuscula, striis rugosis subasperata, diaphana, corneo-rubella ; spira turbi-nata, acuta ; anfr. $3\frac{1}{2}$ convexis, ultimo ventroso, $\frac{2}{3}$ longitudinis formante ; columella leviter arcuata, compressa, albida ; apertura oblique, undique incumbente, subregulariter ovali, intus marg..ri-tacea ; perist. simplice, marginibus subsymmetricis.*

Long. $10\frac{2}{3}$, diam. $6\frac{1}{3}$, alt. 5 mill.

Hab. Flagstaff Hill, St. Helena.

12. **HELIX KERMANDECI**, Pfr. *H. testa imperforata, turbinata, tenui, superne subdistanter striata, hyalina ; spiru convexiusculo-turbinata, subacuminata ; anfr. $5\frac{1}{2}$ -6 convexis, lente accrescenti-bus, ultimo subangulato, antice rotundato ; apertura vix obliqua, lunari ; perist. simplice, recto, marginibus distantibus, columellari superne subreflexo, adnato.*

Diam. maj. $3\frac{2}{3}$, min. $3\frac{1}{3}$, alt. $2\frac{1}{2}$ mill.

Hab. Sunday Island, Kermandee Group (Lieut. Chimmo, R.N.).

13. **HELIX OPHIRIA**, Pfr. *H. testa subperforata, depresse turbinato-globosa, tenuissima, membranacea, radiato-rugata et lineis spirali-bus minutissimis sub lente sculpta, pellucida, virenti-cornea ; spira subconoidea, superne alba ; anfr. 5 convexiusculis, sensim accrescentibus, ultimo peripheria subangulato, antice rotundato, busi inflato ; apertura fere diagonali, rotundato-lunuri ; perist.*

simplice, marginibus dextro et basali subinflexis, columellari fere verticali, superne subreflexo.

Diam. maj. 20, min. 17, alt. $11\frac{1}{2}$ mill.

Hab. Mount Ophir, Malacca (Dr. Traill).

Vitrina heliciformis, Pfr. 1854, is an imperfect form of this species.

14. **HELIX SHIPLAYI**, Pfr. *H. testa perforata, subturbanata, solida; superne arcuato-plicata striisque spiralibus eleganter granulata, isabellina; spira convexiusculo-conica, obtusula; sutura subcanaliculata; anfr. 6 convexis, lente accrescentibus, ultimo peripheria carina acuta, compressa, antice evanescente munito, basi convexo, levicre; apertura diagonali, subangulato-lunari, intus margaritacea; perist. simplice, obtusulo, margine columellari fere verticali, superne triangulatim reflexo.*

Diam. maj. 20, min. 18, alt. $11\frac{1}{2}$ mill.

Hab. Neilgherries, India (Mr. Conway Shipplay).

15. **HELIX ACALLES**, Pfr. *H. testa vix perforata, turbinata, tenui, oblique rugosa, diaphana, pallide cornea; spira conoidea, acutiuscula; anfr. $5\frac{1}{2}$ convexiusculis, sensim accrescentibus, ultimo peripheria carinato, antice non descendente, basi convexiore; apertura obliqua, subangulato-lunari; perist. simplice, recto, margine columellari superne anguste reflexo.*

Diam. maj. 13, min. 11, alt. $7\frac{1}{2}$ mill.

Hab. Neilgherries, India (Mr. Conway Shipplay).

16. **HELIX CALABARICA**, Pfr. *H. testa perforata, turbinato-lentiformi, tenui, levigata, superne liris 6 argutis, filiformibus cincta, diaphana, cornea; spira conoidea, acutiuscula; anfr. 6 convexiusculis, sensim accrescentibus, ultimo non descendente, acute carinato, basi convexo; apertura parum obliqua, angulato-lunari; perist. simplice, recto, marginibus vix convergentibus, columellari arcuato, superne vix reflexo.*

Diam. maj. $9\frac{1}{2}$, min. $8\frac{1}{3}$, alt. $4\frac{1}{2}$ mill.

Hab. Old Calabar, Guinea.

17. **HELIX DARNAUDI**, Pfr. *H. testa perforata, depresso conoideo-globosa, tenui, rugoso-striata, cornea, fasciis opacis albidis notata; spira conoidea; anfr. fere 5 vix convexiusculis, sensim accrescentibus, ultimo non descendente, rotundato; apertura vix obliqua, rotundato-lunari; perist. simplice, recto, marginibus subconvergentibus, columellari superne dilatato, patente.*

Diam. maj. 8, min. 7, alt. 5 mill.

Hab. Sennaar, Interior of Africa (Mr. Darnaud).

18. **HELIX ARGUTA**, Pfr. *H. testa subclause perforata, depressa, tenui, superne argente et confertim arcuato-striata, corneo-fusca; spira parum elevata, obtusa; sutura subcrenulata; anfr. $4\frac{1}{2}$ celester accrescentibus, superis vix convexiusculis, ultimo peripheria carina fusiformi cincto, antice non descendente, basi levior, luteo-virente, medio excavato; apertura diagonali, ampla, lunato-ro-*

tundata, intus margaritacea; perist. simplice, recto, marginibus conniventibus, columellari superne in laminam brevem callosam reflexo.

Diam. maj. 36, min. 29, alt. 18 mill.

Hab. Tenga Hills, Java (C. Shipley, Esq.).

19. **HELIX CHIMMOI**, Pfr. *H. testa umbilicata, convexo-depressa, tenuiuscula, confertim plicato-striata, cornea, rufo irregulariter maculata; spira parum elevata; anfr. 5 convexis, sensim accrescentibus, ultimo non descendentे, subdepresso-rotundato; umbilico conico, $\frac{1}{3}$ diametri subæquante; apertura parum obliqua, rotundato-lunari; perist. simplice, recto, marginibus convergentibus.*

Diam. maj. $3\frac{2}{3}$, min. 3, alt. $1\frac{1}{2}$ mill.

Hab. Sunday Island, Kermandee Group (Lieut. Chimmo, R.N.).

20. **HELIX CONFERTA**, Pfr. *H. testa umbilicata, conoidea, solida, striis incrementi irregularibus et confertissimis spiralibus decussatula, sericea, fulvida; spira conoidea, obtusa; sutura pallide marginata; anfr. vix 5 convexiusculis, sensim accrescentibus, ultimo peripheria angulato, ad suturam turgidulo, basi convexo, juxta umbilicum angustum compresso; apertura perobliqua, subtetragono-lunuri; perist. obtuso, margine supero recto, basali incrassato, reflexiusculo, columellari declivi, subdentato, superne triangulatum reflexo.*

Diam. maj. $34\frac{1}{2}$, min. 30, alt. 20 mill.

Hab. — ?

21. **HELIX DAMAHOYI**, Pfr. *H. testa imperforata, globoso-depressa, tenuiuscula, oblique striata, parum nitente, saturate fusca, fasciis latis nigris, unica lutea mediana et superpositis non-nullis hydrophanis albidis ornata; spira vix elevata, obtusa; anfr. 4 convexiusculis, rapide accrescentibus, ultimo inflato, oblique malleato-plicato; columella declivi, compressa, dilatata, alba; apertura obliqua, truncato-ovali, intus albida; perist. albo, late expanso et reflexiusculo.*

Diam. maj. 47, min. 37, alt. 28 mill.

Hab. Philippine Islands (Mr. Damahoy).

22. **HELIX MEOBAMBENSIS**, Pfr. *H. testa obtecte umbilicata, conoideo-depressa, solida, subdistanter rugoso-striata, undique minute granulata, cinnamomea; spira conoideo-convexa, obtusa; anfr. 5 planiusculis, lente accrescentibus, ultimo antice descendente, peripheria subcarinato, utrinque convextiore; apertura perobliqua, tetragono-lunari; perist. albo, reflexo, marginibus callo junctis, columellari dilatato, adnato, stricto, ad dextram unidentato.*

Diam. maj. 32, min. 26, alt. 15 mill.

Hab. Meobamba, Peru (Mr. Porte).

23. **HELIX AMMIRALIS**, Pfr. *H. testa umbilicata, turbinato-de-*

pressa, solidula, undique subruditer striata, fulva vel lutescente, plerunque fascia unica peripherica nigro-castanea ornata; spira regulariter conoidea; anfr. 6½ vix convexiusculis, lente accrescentibus, ultimo antice vix descendente, peripheria carinato, basi convexo, circa umbilicum angustum, castaneum subcompresso; apertura obliqua, lunari; perist. hepatico, breviter expanso, margine columellari superne fornicatim reflexo.

Diam. maj. 36, min. 33, alt. 20 mill.

Hab. China (Admiral Cecille).

24. **HELIX MUCIDA**, Pfr. *H. testa umbilicata, turbinato-depressa, tenuiuscula, striatula, saturate rufa, quasi mucore obducta; spira conoidea, obtusula; anfr. 5 convexis, sensim accrescentibus, ultimo rotundato, antice descendente, circa umbilicum infundibuliformem subangulata; apertura fere diagonali, rotundato-lunari, intus nitida, carnea; perist. breviter expanso, marginibus vix convergentibus, columellari superne triangulatum dilatato, patente.*

Diam. maj. 20, min. 16½, alt. 11 mill.

Hab. Percy's Island (Lieut. Chimmo, R.N.).

25. **HELIX GUEINZII**, Pfr. *H. testa umbilicata, depressa, carinata, tenui, plicis subdistantibus retrorsum descendantibus sculpta, pellucida, pallide cornea, seriebus macularum rufarum ornata; spira vix elevata; anfr. 5 sensim accrescentibus, prope suturam turgidis, ultimo superne subacute carinato, antice non descendente, subtus per inflato, circa umbilicum ($\frac{1}{6}$ diametri occupantem) conicum levigato, subangulata; apertura vix obliqua, subangulato-lunari; perist. tenui, undique breviter expanso, margine columellari superne dilatato, patente.*

Diam. maj. 20, min. 16, alt. 9 mill.

Hab. Meobamba, Peru (Mr. Gueinzius).

26. **HELIX BASIDENTATA**, Pfr. *H. testa umbilicata, conoideo-subsemiglobosa, solida, oblique striata, alba, fasciis 2 castaneis et 1 angustiore aurantiaca prope suturam ornata; spira convexo-conoidea, obtusa; anfr. 5 convexis, sensim accrescentibus, ultimo carinato, antice perdeflexo, basi planiusculo; umbilico angusto, pervio; apertura horizontali, elliptica; perist. continuo, undique reflexo, margine basali medio dente 1 valido, obtuso armato.*

Diam. maj. 26, min. 20, alt. 11 mill.

Hab. Philippine Islands.

27. **BULIMUS PHÆOSTYLOS**, Pfr. *B. testa imperforata, oblongo-ovata, tenui, oblique confertim striata, pallide fulvescenti-carnea, fasciis 3 nigricantibus (1 suturali, 1 infraperipherica, tertia columellari) ornata; spira convexo-conica, apice obtusula, nigro-violacea; anfr. 5 modice convexis, sensim accrescentibus, ultimo spira paulo breviore; columella compressa, plana, fusca, basi*

subtruncata ; apertura obliqua, truncato-oblonga ; perist. castaneo, anguste reflexo.

Long. 37, diam. 24 mill.

Hab. Philippine Islands.

28. **BULIMUS INÆQUALIS**, Pfr. *B. testa anguste umbilicata, oblique fusiformi, tenuiuscula, lœvigate, nitida, alba unicolo re vel strigis latis ramosis violaceo-nigricantibus picta ; spira gracili, conica, acuta ; anfr. 6 vix convexiusculis, ultimo spira longiore, antice subascendente, latere aperturali planulato, basi subattenuato ; apertura subverticali, elongato-auriformi, intus violaceo-limbata ; columella violacea, superne obliquè plicata ; perist. albo, late expanso, margine columellari plano, patente, superne flexuoso.*

Long. 43, diam. 13 mill.

Hab. Banks of the Maranhon.

29. **BULIMUS GUEINZII**, Pfr. *B. testa profunde rimata, oblongo-conica, tenui, sublœvigate, nitida, albida, strigis angustis fuscis, basin versus undulatis, ornata ; spira conica, acutiuscula ; anfr. 6 convexiusculis, ultimo spira vix breviore, basi subattenuato, ad rimam violaceo ; columella compressa, subtorta, violacea ; apertura vix obliqua, truncato-oblonga, intus lilacina ; perist. tenui, margine dextro superne valde curvato, late expanso, columellari patente.*

Long. 23, diam. 10 mill.

Hab. Meobamba, Peru (*Mr. Gueinzius*).

30. **BULIMUS CLARUS**, Pfr. *B. testa rimato-umbilicata, ovato-oblonga, tenui, lœvigate, nitida, diaphana, sub epidermide fugace fulvescente-albida ; spira elongata, convexo-conica, obtusula ; anfr. 5½ modice convexis, ultimo ⅔ longitudinis subæquante, basi rotundato ; columella leviter arcuata ; apertura obliqua, truncato-ovali ; perist. tenui, marginibus conniventibus, dextro breviter expanso, columellari dilatato, patente.*

Long. 16, diam. 7½ mill.

Hab. Meobamba, Peru (*Mr. Gueinzius*).

31. **BULIMUS LACTIFLUUS**, Pfr. *B. testa breviter rimato-perforata, fusiformi-turrita, tenuiuscula, sublœvigate, cornea, strigis lacteis subserratis notata ; spira turrita, apice acutiuscula, sœpe nigro-cornæa ; anfr. 10-11 convexiusculis, ultimo ⅔ longitudinis subæquante, basi attenuato, compresso ; columella leviter arcuata ; apertura vix obliqua, ovali-oblonga ; perist. tenui, margine dextro anguste expanso, columellari a basi dilatato, fornicatim reflexo.*

Long. 16½-17, diam. 4½ mill.

Hab. Chili.

32. **BULIMUS FLORIDANUS**, Pfr. *B. testa anguste perforata, ovato-turrita, sublœvigate, griseo-hyalina, strigis et maculis opacis, albis notata ; spira elongato-conica, acutiuscula ; anfr. 6½ convexiusculis, superis interrupte fusco-fasciatis, ultimo ⅔ longi-*

tudinis subæquante, infra medium subangulato, basi attenuato; columella subtorta, recedente; apertura parum obliqua, ovali; perist. tenui, margine dextro anguste expanso, columellari dilatato, reflexo, fere adnato.

Long. $15\frac{2}{3}$ –17, diam. $7\frac{1}{2}$ mill.

Hab. Florida.

33. **BULIMUS DISTANS**, Pfr. *B. testa compresse umbilicata, ovato-conica, tenuiuscula, costis subarcuatis, chordæformibus subdistantibus sculpta, subdiaphana, albida; spira elevato-conica, obtusula; anfr. 7 convexis, ultimo $\frac{3}{7}$ longitudinis subæquante, basi rotundato; columella profunde subtorta; apertura parum obliqua, acuminato-subovali; perist. tenui, marginibus convergentibus, dextro breviter expanso, columellari dilatato, patente.*

Long. $22\frac{1}{2}$, diam. $10\frac{1}{2}$ mill.

Hab. Isle of Karah, Gulf of Persia.

34. **BULIMUS ÆSTIVUS**, Pfr. *B. testa umbilicata, oblongo-turrita, solidula, irregulariter striata, albida; spira elongata, convexiusculo-conica, acutiuscula; anfr. 6 modice convexis, ultimo $\frac{3}{7}$ longitudinis subæquante, basi vix attenuato; columella subrecedente; apertura parum obliqua, acuminato-ovali; perist. tenui, margine dextro anguste expanso, columellari dilatato, fornicalim reflexo.*

Long. 17, diam. $7\frac{1}{3}$ mill.

Hab. Meobamba, Peru (Mr. Gueinzius).

35. **BULIMUS CHARBONNIERI**, Pfr. *B. testa profunde rimata, ovato-conica, tenuiuscula, confertim costulato-striata, diaphana, corneo-albida; spira elevato-conica, obtusa; anfr. 6 convexiusculis, ultimo spira paulo breviore, juxta umbilicum rimæformem subcompresso; columella substicta; apertura parum obliqua, elliptico-ovali; perist. tenui, marginibus approximatis, dextro perarcuato, breviter expanso, columellari sursum dilatato, patente.*

Long. 15, diam. 8 mill.

Hab. Isle of Karah, Gulf of Persia.

36. **BULIMUS TIBETANUS**, Pfr. *B. testa profunde rimata, ovato-oblonga, pupæformi, solidula, striatula, albida, strigis et maculis corneis irregulariter notata; spira subcylindrica, sensim in conum obtusulum attenuata; anfr. 8– $8\frac{1}{2}$ vix convexiusculis, ad suturam plicatulis, ultimo $\frac{1}{3}$ longitudinis subæquante, antice ascendentem, busi compresso; columella subuplicata; apertura verticali, sinuato-ovali; perist. albolabiato, expanso, marginibus callo subjunctis, columellari reflexiusculo.*

Long. 32, diam. 11 mill.

Hab. Tibet.

37. **BULIMUS COCHINCHINENSIS**, Pfr. *B. testa imperforata, fusiformi-ovata, solida, lœvigata, nitida, pallide sulphurea vel albida; spira convexiusculo-turrita, apice obtusula; sutura levi, pallida; anfr. 6–7, superis planiusculis, sequentibus convexiori-*

bus, ultimo $\frac{2}{5}$ longitudinis subæquante, basi attenuato, subcompresso; columella uniformi, leviter torta; apertura parum obliqua, elliptico-ovali; perist. subincrassato, marginc dextro breviter expanso, columellari dilatato, adnato.

Long. 39, diam. 17 mill.

Hab. Cochin China.

38. **BULIMUS CHION**, Pfr. *B. testa perforata, oblonga, solida, striatula, alba; spira elongata, in conum acutiusculum terminata; anfr. 7-8 modice convexis, ultimo $\frac{1}{3}$ longitudinis paulo superante, antice subascendente, basi rotundato; apertura verticali, ovato-lunari; perist. calloso, marginibus callo junctis, columellari brevi, substricto, dilatato, patente.*

Long. 12, diam. $5\frac{1}{2}$ mill.

Hab. India; Punjab, Kurrachee, mouth of the Indus (Mr. Conway Shipley).

39. **BULIMUS KANAIENSIS**, Pfr. *B. testa subperforata, conico-ovata, solidula, striatula et irregulariter malleato-impressa, alba; spira conica, sursum interdum grisea, apice obtusa; anfr. 5 convexis, ultimo spiram vix superante, oblique descendente, basi rotundata; columella vix arcuata; apertura obliqua, truncato-ovali; perist. simplice, recto, margine columellari dilatato, reflexo, subappresso.*

Long. 14, diam. 8 mill.

Hab. Kanai, Sandwich Islands.

40. **BULIMUS LORRAINI**, Pfr. *B. testa subperforata, ovato-conica, tenui, rugata, subepidermide fulvida, glutinosa, alba; spira convexiusculo-conica, obtusula; anfr. 4 convexis, ultimo spiram paulo superante, basi parum attenuato; columella compressa, leviter arcuato; apertura obliqua, acuminato-ovali, intus margaritaceo-albida; perist. simplice, recto, margine columellari anguste reflexo, subadnato.*

Long. 20, diam. 9 mill.

Hab. Isle of Penang (Mr. Lorrain).

41. **BULIMUS WOODWARDI**, Pfr. *B. testa perforata, ovato-turrita, solidula, striata et submalleata, nitidula, fulvido-alba; spira elongato-conica, acutiuscula; anfr. 8 convexiusculis, ultimo spira paulo breviore, basi subattenuato; columella recedente; apertura obliqua, ovali-oblonga; perist. simplice, recto, margine dextro leviter arcuato, columellari sursum dilatato, fornicatim reflexo, perforationem angustum semitegente.*

Long. 31, diam. $13\frac{1}{2}$ mill.

Hab. Andes of Peru.

42. **BULIMUS CASTELNEAUI**, Pfr. *B. testa subperforata, fusiformi-turrita, tenuiuscula, striatula (sub lente decussatula), nitida, griseo-albida, punctis rarissimis pellucidis conspersa; spira elongato-conica, apice acuta, cornea; anfr. $7\frac{1}{2}$ convexiusculis, ultimo $\frac{2}{5}$ lon-*

gitudinis subæquante, basi attenuato; columella substricta; apertura obliqua, oblonga, intus carnea; perist. simplice, recto, margine columellari sursum dilatato, fornicato-reflexo.

Long. 20, diam. $8\frac{1}{2}$ mill.

Hab. Rio Pampas, Bolivia (Mr. Castelneau).

43. **BULIMUS NIGROAPICATUS**, Pfr. *B. testa perforata, ovato-conica, tenuiuscula, striata, nitida, albida, fasciis nigricantibus crebris, superioribus moniliformibus, ornata; spira conica, apice acutiuscula, nigra; anfr. $5\frac{1}{2}$ parum convexis, ultimo spira paulo langiore, ventroso; columella levissime arcuata; apertura obliqua, elliptico-ovali; perist. simplice, recto, murgine columellari superne late reflexo, subappresso.*

Long. 22, diam. $11\frac{1}{2}$ mill.

Hab. Rio Pampas, Bolivia (Mr. Castelneau).

44. **BULIMUS STENACME**, Pfr. *B. testa umbilicata, ovato-turrita, solidula, ruguloso-striata, albida, strigis angustis pallide corneis variegata; spira elongata, apicem versus acutiusculum attenuata; anfr. 7 convexiusculis, ultimo $\frac{2}{5}$ longitudinis subæquante, juxta umbilicum angustum subcompresso; apertura obliqua, oblongo-ovali, intus fusco-carnea; perist. simplice, recto, margine columellari superne dilatato, fornicatum reflexo.*

Long. $20\frac{1}{2}$, diam. 9 mill.

Hab. Bolivia.

45. **BULIMUS MONACHUS**, Pfr. *B. testa anguste perforata, oblongo-turrita, tenui, striatula, diaphana, sordide cornea; spira convexo-turrita, apice acutiuscula; anfr. $7\frac{1}{2}$ vix convexiusculis, ultimo $\frac{3}{4}$ longitudinis subæquante, basi parum attenuato, rotundato; columella recedente; apertura obliqua; oblongo-ovali; perist. simplice, recto, margine columellari sursum dilatato, reflexo.*

Long. 31, diam. $11\frac{1}{2}$ mill.

Hab. Meobamba, Peru (Mr. Gueinzii).

46. **PARTULA PURPURASCENS**, Pfr. *P. testa perforata, ovato-conica, solidu, oblique striata striisque spiralibus confertis distinete decussata, nitida, purpurascenti-fusca; spira convexo-conica, acuta; sutura levi; anfr. $5\frac{1}{2}$ vix convexiusculis, ultimo spiram subæquante, peripheria subangulato, basi rotundato; columella leviter arcuata; apertura parum obliqua, truncato-oblonga; perist. fusco-violaceo, undique patente et reflexiusculo.*

Long. 22, diam. 14 mill.

Hab. —?

47. **PARTULA CALLIFERA**, Pfr. *P. testa umbilicata, ovato-conica, solida, sublævigata (sub lente punctulato-striata), albida; spira convexiusculo-conica, obtusula; sutura levi; anfr. 5 convexiusculis, ultimo spira paulo longiore, inflato; columella subverticali, superne tuberculifera, intus plicata; apertura vix obliqua, sinuato-oblonga, dente profundo parietali coarctata; perist. incrassato,*

dilatato, marginibus callo junctis, dextro supra medium callo oblongo intus munito.

Long. 19, diam. 11 mill.

Hab. — ?

48. PARTULA LÆVIGATA, Pfr. *P. testa profunde rimata, ovato-conica, solida, lævigata (sub lente vix striatula), nitida, lutescente; spira conica, obtusula; sutura mediocri; anfr. 5 convexiusculis, ultimo spiram subæquante, prope sutaram tumidiore, basi rotundato; columella supra medium introrsum nodoso-plicata; apertura vix obliqua, oblonga, tuberculo profundo anfractus penultimi coarctato; perist. crasso, albo, undique patente, margine columellari superne dilatato, adnato.*

Long. 20, diam. 10 mill.

Hab. — ?

49. PARTULA LILACINA, Pfr. *P. testa subobtecte perforata, ovato-conica, solidu, sub lente spiraliter striata, nitidula, lilacina; spira conica, acuta; sutura levi; unfr. $5\frac{1}{2}$ vix convexiusculis, ultimo spiram subæquante, rotundato; columella superne tuberculo subcircumscripto munita; apertura obliqua, truncato-oblonga; perist. incrassato, albido, breviter expanso, margine columellari fere adnato.*

Long. $17\frac{1}{2}$, diam. 10 mill.

Hab. Marquesas Islands.

50. ACHATINELLA (ACHATINELLASTRUM) OVUM, Pfr. *A. testa sinistrorsu, imperforata, globoso-conica, solida, rugulosa-striata, nitidala, albida; spira concaviusculo-conica, apice acuta; sutura profunde marginata; anfr. $5\frac{1}{2}$, superis planis, sequentibus convexis, ultimo inflato, spira vix breviore; plica columellari crassa, tuberculiformi, pallide lilacea; apertura diagonali, sinuato-semicirculari; perist. recto, nigro-fusco, limbato, intus crenato-labiato.*

Long. $19\frac{1}{2}$, diam. 13 mill.

Hab. Oahu, Sandwich Islands (Dr. Newcomb).

51. ACHATINELLA (LAMINELLA) FARCIMEN, Pfr. *A. testa sinistrorsa, subperforata, oblongo-conica, solidula, rugulosa, sub epidermide nigricante albida; spira superne in conum acutum nudum terminata; anfr. 7, superis planis, 2 ultimis perconvexis, ultimo $\frac{1}{3}$ longitudinis paulo superante, medio subangulato; plica columellari compressa, obliqua; apertura obliqua, semiovuli, intus alba; perist. simplice, recto, margine columellari subpatente.*

Long. 19, diam. 9 mill.

Hab. Mani, Sandwich Islands (Dr. Newcomb).

52. ACHATINELLA (LABIELLA) CALLOSA, Pfr. *A. testa imperforata, dextrorsa, fusiformi-oblonga, solida, leviter striatula, sub epidermide tenui fulvida alba; spiru elongata, ventroso-conica,*

13m

apice obtusula; sutura sublacerata; anfr. 8 planiusculis, ultimo $\frac{1}{3}$ longitudinis paulo superante, basi attenuato; plica columellari acute dentiformi, alba; apertura vix obliqua, acuminato-elliptica; perist. calloso, obtuso, margine dextro intus obsolete dentato.
 Long. 16, diam. 6 mill.
Hab. Oahu, Sandwich Islands (Dr. Newcomb).

53. *ACHATINA (VARICELLA) GUADELOUPENSIS*, Pfr. *A. testa oblongo-turrata, texui, laevigata, nitidissima, pellucida, virenticornea, varicibus arcuatis, vix prominulis, castaneo-marginatis passim munita; spira regulariter attenuata, apice obtusa; anfr. 7 convexiusculis, ultimo $\frac{1}{3}$ longitudinis subæquante, basi attenuato; columella perarcuata, basi anguste truncata; apertura vix obliqua, acuminato-ovali; perist. tenui, margine dextro antrorum dilatato, rufo-limbato.*

Long. 14, diam. $4\frac{1}{3}$ mill.

Hab. In insula Guadeloupe (Mr. Caillet).

54. *SPIRAXIS SANDWICHENSIS*, Pfr. *Sp. testa subperforata, oblongo-turrata, solidula, cerea; spira turrita, obtusula; anfr. 7 $\frac{1}{2}$ planiusculis, infra suturam plicatis, ultimo $\frac{1}{3}$ longitudinis paulo superante; columella compressa, torta; apertura vix obliqua, ovali; perist. simplice, marginibus callo tenui junctis, dextro antrorum subdilatato, columellari subreflexo.*

Long. 9, diam. 3 mill.

Hab. Sandwich Islands (Dr. Newcomb).

55. *SPIRAXIS OBSOLETA*, Pfr. *Sp. testa subperforata, oblongo-turrata, tenui, levissime striatula, pellucida, nitida, pallide cornea; spira convexiusculo-turrata, obtusa; sutura marginata; anfr. 6 $\frac{1}{2}$ convexiusculis, ultimo $\frac{2}{5}$ longitudinis formante; columella acuta, medio leviter torta; apertura vix obliqua, elliptico-ovali; perist. simplice, margine columellari brevissime reflexo, subadnato.*

Long. 8, diam. 3 mill.

Hab. Sandwich Islands (Dr. Newcomb),

56. *TORNATELLINA GOULDII*, Pfr. *T. testa ovato-conica, tenui, sublaevigata, pellucida, cornea; spira elongato-conica, apice obtusula; anfr. 5 $\frac{1}{2}$ convexis, ultimo $\frac{2}{5}$ longitudinis subæquante, rotundata; apertura obliqua, erecto-lunari, lamella valida intrante parietali et plica crassa triangulari columellæ coarctata; perist. recto, tenui.*

Long. 4, diam. $2\frac{1}{2}$ mill.

Hab. —?

57. *TORNATELLINA NEWCOMBI*, Pfr. *T. testa perforata, ovato-turrata, tenui, sublaevigata, parum nitida, diaphana, pallide cornea; spira subrectilineari, conica, obtusula; anfr. 7 subplanis, ultimo $\frac{1}{3}$ longitudinis vix formante, basi conveexo; apertura obliqua, ro-*

tundato-lunari, lamella mediocri parietali et plicis 2 parallelis columellæ coarctata; perist. simplice, recto.

Long. 4, diam. $2\frac{1}{3}$ mill.

Hab. Sandwich Islands (Dr. Newcomb).

58. **TORNATELLINA PERFORATA**, Pfr. *T. testa perforata, ovato-turrita, arcuato-striata, tenui, albido-hyalina; spira elongata, apice acuta; anfr. 7 convexis, ad suturam distanter plicatulis, ultimo $\frac{1}{3}$ longitudinis paulo superunte; lamella parietali minuta; apertura vix obliqua, sinuato-ovali, acuminata; columella lamina torta munita; perist. tenui, margine dextro recto, superne antrorsum arcuato, basali expanso, columellari fornicatim reflexo, patente.*

Long. 18, diam. $7\frac{1}{2}$ mill.

Hab. Venezuela.

6. DESCRIPTIONS OF SIXTEEN NEW SPECIES OF PNEUMONOPOMA, FROM THE COLLECTION OF H. CUMING, ESQ.

BY DR. L. PFEIFFER.

A. OPISOPHTHALMA.

1. **DIPLOMMATINA CANTORI**, Pfr. *D. testa subrimata, sinistrorsa, ovato-oblonga, tenuiuscula, confertim et oblique plicata, albida; spira ovata, in conum acutiusculum terminata; anfr. 6 convexiusculis, ultimo ascendentem, $\frac{1}{3}$ longitudinis vix superante; apertura subverticali, circulari; perist. subsimplice, continuo, superne adnato, cæterum breviter expanso.*

Long. 2, diam. 1 mill.

Hab. Lord Howe's Island, New Hebrides (Mr. Maegillivray).

2. **TRUNCATELLA CEYLANICA**, Pfr. *Tr. testa subrimata, attenuato-cylindracea, striatula, pellucida, nitida, corneo-rufa vel flavescente; anfr. superst. 4, supremis 2 rotundatis, reliquis planioribus, omnibus ad suturam plicato-crenatis, ultimo basi non compresso; apertura verticali, ampla, angulato-subovali; perist. continuo, margine dextro tenui, expansiusculo, columellari leviter arcuato, apppresso.*

Long. 6, diam. 2 mill.

Hab. Ceylon.

3. **TRUNCATELLA TERES**, Pfr. *Tr. testa vix subrimata, cylindrica, tenuiuscula, longitudinaliter magis minusve distincte costulata, pellucida, nitida, rufo-cornea; sutura marginata, valide plicato-crenata; anfr. superst. 4 subæqualibus, convexiusculis, ultimo basi breviter cristato, calloso, albido; apertura verticali, late orali, superne angulata, ad dextram dilatata; perist. simplice, continuo, margine dextro expansiusculo, columellari adnato.*

Long. 6, diam. 2 mill.

Hab. Isle of Mauritius, and Trinity Bay, Australia.

4. **TRUNCATELLA BARBADENSIS**, Pfr. *Tr. testa subrimata, cylindracea, sursum vix attenuata, solidula, costulis confertis, obtusis, subrectis regulariter sculpta, sericina, rufo-cornea; sutura profunda; anfr. superst. 4½ perconvexis, lente accrescentibus, ultimo basi crista albida, antice peristoma cingente munito; apertura verticali, ovali, superne subrotundata, basi subeffusa; perist. continuo, margine dextro expanso et reflexiusculo, perarcuato, columellaris subadnato.*

Long. 6½, diam. 2 mill.

Hab. Island of Barbadoes, West Indies.

B. ECTOPHTHALMA.

5. **CYCLOSTOMA (CYCLOTUS) DAUCINUM**, Pfr. *C. testa umbilicata, depressa, solidula, subangulata, daucina vel albida; spira parum elevata; sutura simplice; anfr. 4½ convexis, sensim accrescentibus, superioribus spiraliter striatis, ultimo rugoso, carina mediocri, antice evanescente munito; umbilico conico, ¼ diametri paulo superante; apertura parum obliqua, subcirculari; perist. simplice, recto, ad anfr. penultimum breviter interrupto. Operc. anguste et obsolete spiratum.*

Diam. maj. 12, min. 10, alt. 6 mill.

Hab. Salomon's Islands.

6. **CYCLOSTOMA (OPISTHOPORUS) COCHINCHINENSE**, Pfr. *C. testa late umbilicata, discoidea, solida, vix striatula, sub epidermide fulvida alba, radiis pellucidis notata; spira subplana, medio vix elevata; anfr. 5 convexis, ultimo terete, antice soluto, pone aperturam spiraculo versus anfr. penultimum currato munito; apertura circulari; perist. subsimplice, vix expansiusculo, superne levissime inciso. Operc. ?*

Diam. maj. 20, min. 16, alt. 7 mill.

Hab. Cochin China.

7. **CYCLOSTOMA (OPISTHOPORUS) EURYOMPHALUM**, Pfr. *C. testa late umbilicata, deppressa, subdiscoidea, solidula, striatula, lutescente, strigis angulosis castaneis, superne latis, subtus linearibus, picta; spira plana, vertice nigro vix prominulo; anfr. 4½, prope suturam canaliculatam subangulatis, ultimo antice vix descendente, 4 mill. pone aperturam spiraculo brevi retroflexo munito; umbilico dimidium diametri fere occupante; apertura diagonali, subcirculari; perist. duplice, interno breviter expanso, ad anfr. penultimum subinciso, externo superne alatim dilatato, latere dextro patente, sinistro obsoleto. Operc. calcareum, angustispirum.*

Diam. maj. 15, min. 12, alt. 4 mill.

Hab. Borneo.

8. **CYCLOSTOMA (CYCLOPHORUS) SHIPLAYI**, Pfr. *C. testa umbilicata, deppressa, tenuiuscula, membranaceo-striata, fulva, cas-*
No. CCCXXII.—PROCEEDINGS OF THE ZOOLOGICAL SOCIETY

taneo oblique strigata; spira vix eleata; anfr. 4 convexis, ultimo terete; umbilico $\frac{1}{3}$ diametri occupante; apertura obliqua, subcirculari; perist. simple, recto, ad anfractum contingum vix interrupto. Operc.?

Diam. maj. 7, min. 6, alt. 3 mill.

Hab. Neilgherries, India (Mr. Conway Shipley).

9. CYCLOSTOMA (LEPTOPOMA) SIGNATUM, Pfr. *C. testa perforata, globoso-turbinata, tenui, sub lente decussatula et lineis filaribus, subelevatis, subdistantibus cincta, diaphana, lutescenti-cornea, strigis confertis rufis fulgurata; spira turbinata, acutiuscula; anfr. 5 convexis, ultimo spiram subaequante, 5-lirato; apertura diagonali, subcirculari; perist. subincrassato, patente, marginibus fere contiguis, columellari angustiore. Operc.?*

Diam. maj. 11, min. $8\frac{2}{3}$, alt. $8\frac{2}{3}$ mill.

Hab. Borneo.

10. CYCLOSTOMA (LEPTOPOMA) DUPLICATUM, Pfr. *C. testa anguste umbilicata, globoso-turbinata, solidula, undique confermit spiraliter striata lirisque filiformibus distantibus cincta, fulva, strigis fulguratis rufis ornata; spira elevata, acutiuscula; anfr. 5 convexis, ultimo superne turgido, infra medium carina levi munito; apertura obliqua, subangulato-circulari; perist. subcontinuo, albo, duplice, interno rix porrecto, externo expanso et reflexinsculo, latere sinistro quasi absciso. Operc.?*

Diam. maj. 10, min. 8, alt. 7 mill.

Hab. —?

11. CYCLOSTOMA (CYCLOSTOMUS) BOIVINI, Pfr. *C. testa obtecta perforata, globoso-turbinata, tenuiuscula, spiraliter obsolete lirata; griseo et fulrido variegata, spadiceo multifasciata; spira turbinata; anfr. fere 6 turgidis, celeriter accrescentibus, ultimo ventroso, peripheria carina 1 compressa, albida munito, circa umbilicum confertim spiraliter lirato; apertura subverticalli, oblongo-rotundata, intus nigricanti-sanguinea; perist. tenui, ad anfr. penultimum subemarginato, superne producto, margine dextro et basali late patente, columellari angusto, supra umbilicum dilatato, adnato. Operc.?*

Diam. maj. 29, min. 23, alt. 25 mill.

Hab. Nos-bé, Madagascar (Mr. Boivin).

12. CYCLOSTOMA (CYCLOSTOMUS) MICROCHASMA, Pfr. *C. testa umbilicata, depresso turbinato-globosa, tenuiuscula, spiraliter confertim lirata, striis incrementi vix decussatula, albido-lutescente; spira turbinata, obtusula; anfr. $4\frac{1}{2}$ convexis, ultimo terete, infra peripheriam fascia latiuscula castanea ornato; umbilico conico, $\frac{1}{3}$ diametri fere occupante; apertura parum obliqua, parvula, fere circulari; perist. tenui, vix expansius-*

culo, marginibus fere contiguis, ad anfr. penultimum callo brevi junctis. Operc.?

Diam. maj. 18, min. $15\frac{1}{2}$, alt. 12 mill.

Hab. Madagascar.

13. **CYCLOSTOMA (CYCLOSTOMUS) SARCODES**, Pfr. *C. testa umbilicata, turbinata, solidula, liris obtusis subconfertis striisque illas transgredientibus creberrimis sculpta, carnea, fusco-violaceo uni- vel plurifasciata; spira turbinata, acutiuscula; anfr. 5 convexis, ultimo circa umbilicum angustum, subpervium liris angustioribus, prominentioribus munito; apertura fere verticali, ovali-rotundata; perist. anguste expanso, breviter adnato, superne subangulato. Operc.?*

Diam. maj. 17, min. 14, alt. 14 mill.

Hab. Madagascar.

14. **CATAULUS CUMINGI**, Pfr. *C. testa vix rimata, turrito-fusiformi, solidula, subgranulato-striata, daucino-fusca; spira convexo-turrita, apice acutiuscula; sutura submarginata; anfr. 8 vix convexiusculis, penultimo convexiore, ultimo attenuato, basi axin vix excedente; carina umbilicali compressa, albida, angulatim patula; periomphalo mediocri, turgido, distinctius striato; apertura subcirculari; perist. albo, continuo, incrassato, fornicatim patente, superne anguste adnato, cæterum dilatato, basi longe producto, canali infundibuliformi perforato.*

Long. $20\frac{2}{3}$, diam. medio 7 mill.

Hab. Ceylon (Mr. Thwaites).

15. **HELICINA IDEÆ**, Pfr. *H. testa conoideo-depressa, tenuiuscula, striata, lineis spiralibus impressis distantibus sculpta, nitida, lutea vel carnea; spira conoidea, acutiuscula; anfr. 5 planiusculis, ultimo peripheria subangulato, basi convexiore; callo basali tenui, concolore; columella brevi, antrorsum subdentata; apertura diagonali, subsemicirculari; perist. acuto, breviter expanso, intus labiato. Operc.?*

Diam. maj. 8, min. $6\frac{1}{2}$, alt. $4\frac{1}{2}$ mill.

Hab. Ceram (Mrs. Ida Pfeiffer).

16. **HELICINA VIRENS**, Pfr. *H. testa conoidea, solida, striatula et sub lente minute spiraliter striata, carinata, nitida, albido-virente; spira convexo-conoidea, apice acuta; sutura marginata, carina interdum prominente; anfr. 6 vix convexiusculis, summis interdum rufo-fasciatis, ultimo antice vix descendente, peripheria acute albocarinato, basi parum convexo; apertura diagonali, subtriangulari-semiovali; columella brevi, superne leviter impressa, callum emitte circumscripturn, luteum, subgranulatum; perist. acuto, albo, expanso. Operc. tenue, castaneum.*

Diam. maj. $11\frac{1}{3}$, min. 10, alt. 8 mill.

Hab. —?