

PROCEEDINGS

OF THE

ZOOLOGICAL SOCIETY

OF LONDON.



PART XXIV.

1856.

PRINTED FOR THE SOCIETY ;
SOLD AT THEIR HOUSE IN HANOVER-SQUARE,
AND BY MESSRS. LONGMAN, BROWN, GREEN AND LONGMANS,
PATERNOSTER-ROW.

tura obliqua, subangulato-circulari; perist. subduplice interno vix interrupto, adnato, externo patente, concentricè striatulo, latere sinistro dilatato, fornicatim reflexo.—Operc. planum, fulvum.

Diam. maj. 12, min. 10, alt. 10 mill.

17. *HELICINA GRATIOSA*, Pfr. *H. testa subgloboso-turbinata, tenuiuscula, striatula, striolis spiralibus obsoletissime notata, nitida, pellucida, fusciscenti-rubra vel lutea; spira conoidea, acutiuscula; anfr. 5 convexiusculis, ultimo subcarinato, antice rotundato; columella subecedente, alba, callum emittente albidum; apertura obliqua, subtriangulato-semicirculari; perist. undique breviter expanso.*—Operc. tenue, solidum, curvum.

Diam. maj. 9, min. $7\frac{1}{2}$, alt. fere 6 mill.

18. *HELICINA SUAVIS*, Pfr. *H. testa turbinata, tenui, nitidissima, superne striatula striisque spiralibus conferte notata, rubella, albido-marmorata, vel lutea aut carnea, pallide rubro variegata; spira conoidea, acuta; anfr. fere 5 convexiusculis, ultimo compresse et acute carinato, basi convexiore; columella brevi, basi subdentata, callum emittente crassiusculum diffusum; apertura obliqua, subtriangulari; perist. albo, margine supero breviter expanso, basali reflexiusculo.*—Operc.?

Diam. maj. $10\frac{1}{2}$, min. $8\frac{3}{4}$, alt. $6\frac{2}{3}$ mill.

Besides the described new species, there were in the same collection from the Admiralty Islands, *Helix Novæ Hiberniæ*, Quoy, *fringilla*, Pfr. (beautiful and large varieties), *migratoria*, Pfr., *Sachalensis*, Pfr., *motacilla*, Pfr., *Cleryi*, Recl., *helicinoides*, Jacq., *pyxis*, Hinds, *Grimardi*, Desh., *brevipila*, Pfr., *Pfeifferi*, Phil., &c.

5. DESCRIPTIONS OF THIRTY-THREE NEW SPECIES OF LAND-SHELLS, FROM THE COLLECTION OF H. CUMING, ESQ.

By DR. L. PFEIFFER.

(Mollusca, Pl. XXXVI.)

1. *HELIX GLORIOSA*, Pfr. *H. testa imperforata, depresso ovatu, crassa, ponderosa, oblique conferte plicato-striata lirisque crebris obtusis subregularibus cincta, fulvida, strigis singulis pallidis et saturatoribus notata; spira convexa, obtusa, nuda, sub lente granulata, apice subtilissimo; anfr. $3\frac{1}{2}$ rapide accrescentibus, ultimo perinflato; apertura diagonali, lunato-ovali, intus rubella; perist. crassissimo, breviter expanso, marginibus callo crasso junctis, columellari adnato.*

Diam. maj. 64, min. 50, alt. 39 mill.

Hab. Madagascar.

2. *HELIX DUCTILIS*, Pfr. *H. testa umbilicata, discoidea, tenui,*
No. CCCXXV.—PROCEEDINGS OF THE ZOOLOGICAL SOCIETY.

sub lente confertim striatula, albido-hyalina; spira plana; anfr. fere 5 subplanis, sensim accrescentibus, ultimo non descendente, depresso, peripheria rotundato, basi vix convexiore; umbilico perspectivo, $\frac{1}{2}$ diametri superante; apertura obliqua, late lunari; perist. simplice, recto, marginibus vix convergentibus, columellari brevi, verticali, cum basali subangulatim juncto.

Diam. maj. $6\frac{1}{2}$, min. $5\frac{1}{2}$, alt. 2 mill.

Hab. Drayton Range, North Australia (Mr. Stutchbury).

3. *HELIX STUTCHBURYI*, Pfr. *H. testa sub obtecte perforata, turbinato-globosa, tenui, superne sub lente minute granulata, pallide fulva, ad suturam et supra medium rufo-fasciata; spira convexo-conoidea, obtusula; anfr. 5 convexiusculis, ultimo rotundato, antice vix descendente, basi lævigato; apertura diagonali, rotundato-lunari; perist. tenui, marginibus subconvergentibus, dextro vix expansiusculo, basali breviter reflexo, columellari fornicatim dilatato, umbilicum angustissimum fere tegente.*

Diam. maj. $15\frac{1}{2}$, min. $13\frac{1}{2}$, alt. 10 mill.

Hab. Drayton Range, North Australia (Mr. Stutchbury).

4. *HELIX DELTA*, Pfr. *H. testa sub obtecte perforata, trochiformi, tenuiuscula, striatula et foveolata, opaca, carneo-albida; spira regulariter conica, apice acuta; anfr. 6 planis, ultimo non descendente, subacute carinato, basi convexiusculo; apertura fere diagonali, angulato-lunari; perist. simplice, margine dextro recto, basali reflexiusculo, columellari supra perforationem dilatato, patente.*

Diam. maj. 8, min. $7\frac{1}{2}$, alt. $6\frac{1}{2}$ mill.

Hab. Drayton Range, North Australia (Mr. Stutchbury).

5. *HELIX Pliculosa*, Pfr. *H. testa umbilicata, turbinato-globulosa, tenuiuscula, superne confertim plicosa, diaphana, parum nitida, rubello-cornea; spira conoidea, vertice subtili; anfr. $4\frac{1}{2}$ convexiusculis, ultimo supra medium subangulato, antice rotundato, vix descendente, basi inflato, circa umbilicum angustum, pervium subcompresso; apertura obliqua, lunato-rotundata; perist. albido, marginibus subconniventibus, dextro expansiusculo, columellari sursum dilatato, cum basali angulum obtusum formante.*

Diam. maj. 16, min. 13, alt. 10 mill.

Hab. Drayton Range, North Australia (Mr. Stutchbury).

6. *HELIX CASTRENSIS*, Pfr. *H. testa imperforata, conoideo-semiglobosa, solida, rugoso-plicatula et irregulariter subgranulata, pallide fulvida, strigis fuscis crebris regulariter radiata; spira convexiusculo-conoidea, obtusula; anfr. 4 vix convexiusculis, ultimo medio subacute carinato, antice breviter descendente, inflato, basi pallidior; apertura perobliqua, subtriangulari-lunari; perist. albo, margine dextro arcuato, breviter reflexo, basali dilatato, adnato, introrsum dentibus 2 obtusis plicæformibus munito.*

Diam. maj. 25, min. 21, alt. 15 mill.

Hab. West Indies.

7. **HELIX LINDSTEDTI**, Pfr. *H. testa sinistrorsa, perforata, conoideo-leuticulari, carinata, tenui, diaphana, superne undulato-striata et minutissime granulata, parum nitida, rubello-lutescente; spira regulariter elevata, obtusa; sutura lineari; anfr. 6 planiusculis, sensim accrescentibus, ultimo non descendente, supra medium acute carinato, basi convexo, nitidissimo, non decussato, circa perforationem non apertam albido; apertura fere diagonali, securiformi; perist. simplice, recto, margine columellari in laminam brevem revolutam dilatato.*
 Diam. maj. $35\frac{1}{2}$, min. 31, alt. 14 mill.
Hab. Malacca (Rev. F. W. Lindstedt).
8. **HELIX BOURGUIGNATI**, Pfr. *H. testa umbilicata, globoso-depressa, tenui, minutissime malleato-striata, parum nitente, diaphana, carnea; spira breviter conoidea, vertice subtili, luteo; anfr. 6 convexiusculis, regulariter accrescentibus, ultimo rotundato, antice vix descendente; umbilico angusto, vix pervio; apertura obliqua, rotundato-lunari; perist. intus albolabiato, marginibus vix convergentibus, dextro recto, basali reflexiusculo, ad umbilicum dilatato, patulo.*
 Diam. maj. 19, min. 16, alt. 11 mill.
Hab. Crimea.
9. **HELIX ADELAIDÆ**, Pfr. *H. testa umbilicata, depressa, carinata, tenuiuscula, striata, alba, fascia fuscata supera ornata; spira vix elevata, vertice obtusiusculo, corneo; anfr. 4 convexiusculis, ultimo non descendente, supra peripheriam subacute carinato, basi inflato; umbilico perspectivo, $\frac{1}{4}$ diametri fere æquante; apertura fere diagonali, lunato-rotundata, ad carinam vix angulata; perist. recto, intus levissime labiato, marginibus conniventibus, columellari subpatulo.*
 Diam. maj. 6, min. 5, alt. fere 3 mill.
Hab. Adelaide, Australia.
10. **HELIX CHIONODISCUS**, Pfr. *H. testa subobtecte umbilicata, conoideo-lentiformi, carinata, solida, rugoso-striatula, opaca, nivea; spira convexiusculo-conoidea, nucleo læviusculo, obtuso; anfr. 5 vix convexiusculis, ultimo antice breviter deflexo, infra suturam linearem turgidulo, periphæria magis minusve acute carinato, basi convexo; apertura diagonali, rhombeo-lunari; perist. subcalloso, marginibus subparallelis, dextro recto, basali perarcuato, reflexo, versus umbilicum sensim dilatato, lamina lata adnata illum fere claudente.*
 Diam. maj. 26, min. 23, alt. 13 mill.
Hab. Crimea.
11. **HELIX GREVILLEI**, Pfr. (Pl. XXXVI. fig. 8.) *H. testa imperforata, conica, solida, striatula et plicis validis oblique antrorsum descendentibus, interdum malleato-interruptis munita, castanea, strigis sparsis nigricantibus notata; spira conoidea,*

apice rosea, acutiuscula; anfr. 4½ convexiusculis, rapide accrescentibus, penultimo subtus albo-callosa, ultimo antice descendente, peripheria obtuse sed distincte angulato; apertura perobliqua, truncato-oblonga, intus pallide cærulescente; perist. nigro, expanso et reflexo, marginibus parallelis, callo nigro junctis, columellari dilatato, plano, adnato, dextrorsum subdentato, ad axin albido.

Diam. maj. 49–58, min. 36–42, alt. 32–39 mill.

Hab. Ceylon (*Mr. Thwaites*).

12. *ENNEA INSIGNIS*, Pfr. *E. testa breviter rimata, ovato-oblonga, tenuissima, oblique confertim striata, striisque spiralibus versus basin evanescentibus decussatula, nitida, pellucida, cerco-hyalina; spira ovoidea, apice obtusa; sutura submarginata; anfr. 7 convexiusculis, penultimo supra aperturam subplanato, ultimo ¾ longitudinis formante, antice arcuatim ascendente, juxta rimam subcompressa; columella recedente, dentato-plicata; perist. tenui, albo, undique expanso, murgine columellari dilatato, patente.*

Long. 37, diam. 20 mill.

Hab. Gaboon, Africa (*Mr. Auboy*).

13. *BULIMUS PALAVANENSIS*, Pfr. *B. testa imperforata, oblonga, solida, striatula, epidermide fusco-cinerea, castaneo varie strigata, sursum detrita munita; spira convexiusculo-conica, apice obtusula; anfr. 5–5½ modice convexis, ultimo spiram subæquante vel brevior; columella substricta, albida; apertura obliqua, truncato-ovali, intus griseo-cærulescente; perist. callosa, nigricante, breviter reflexo.*

Long. 48–49, diam. 27 mill.

Hab. Palawan (*Dr. Traill*).

14. *BULIMUS LIBROSUS*, Pfr. *B. testa imperforata, ovato-oblonga, solida, striatula, epidermide subhydrophana, libro-cinerea, fusco irregulariter strigata et ad suturam ocellata obducta; spira convexiusculo-conica, obtusa; anfr. 5½ planiusculis, ultimo spira vix brevior, epidermide decidua circa columellam pallidam, compressam, substrictam saturate castanea, nitida; apertura obliqua, ovali oblonga, intus cærulescente; perist. castaneo-nigro, subincrassato, revoluto, margine dextro leviter arcuato.*

Long. 40, diam. 21 mill.

Hab. Palawan (*Dr. Traill*).

15. *BULIMUS LINDSTEDTI*, Pfr. *B. testa sinistrorsa, imperforata, ovato-conica, solida, striatula, nitida, candida; spira elongato-conica, apice obtusula; anfr. 6½ convexiusculis, ultimo ¾ longitudinis subæquante, obsolete angulato, antice rotundato; columella subverticali, vix torta; apertura fere diagonali, subsemicirculari; perist. leviter incrassato, expanso, marginibus callo concolore junctis.*

Long. 39, diam. 17 mill.

Hab. Malacca (*Rev. F. W. Lindstedt*).

16. *BULIMUS PARALLELUS*, Pfr. *B. testa compressa umbilicata, fusiformi, tenuiuscula, striatula, fulva, epidermide albida subreticulata; spira elongato-conica, obtusa; anfr. 7 modice convexis, ultimo spira paulo brevior, basi compresso; columella superne leviter torto-plicata; apertura verticali, oblonga, lateribus parallelis, basi effusa, utrinque fusco-strigata; perist. tenui, albo, expanso, margine dextro supra medium impresso, columellari latiore, sulco arcuato ab anfractu contiguo separato.*
 Long. 22, diam. 7 mill.
Hab. St. Catherine's, Brazil.
17. *BULIMUS CATHARINÆ*, Pfr. *B. testa breviter rimata, subperforata, fusiformi, solidula, undique leviter punctato-rugulosa, albida, punctis corneis substrigatim conspersa; spira ventroso-turrita, apice acutiuscula; anfr. $8\frac{1}{2}$ convexiusculis, ultimo $\frac{1}{3}$ longitudinis vix attingente, basi crista valida, compressa et pone illam minore munito, antice interne et externe interrupte nigro-strigato; apertura obliqua, subseptemdentata; plica 1 lamellæformi in pariete, secunda subquadrangulæ ad columellam, tertia obliqua in latere sinistro baseos effusa, 3 subæqualibus et 1 minuta in margine dextro; perist. albo, undique modice expanso.*
 Long. 23-25, diam. $7\frac{1}{2}$ -8 mill.
Hab. St. Catherine's, Brazil.
18. *BULIMUS SUGILLATUS*, Pfr. *B. testa anguste umbilicata, oblongo-turrita, tenui, irregulariter plicato-striata, cornea, strigis subpunctatis opacis albis, singulisque rufis notata; spira elongato-conica, acuta; sutura levissime crenulata; anfr. 9 convexis, ultimo $\frac{2}{3}$ longitudinis subæquante, basi subcompresso; columella leviter et stricte recedente; apertura parum obliqua, oblonga; perist. simplice, margine dextro recto, columellari sursum dilatato, fornicatim reflexo.*
 Long. 24, diam. $9\frac{1}{2}$ mill.
Hab. Bolivia.
19. *BULIMUS GAYI*, Pfr. *B. testa subobtecte umbilicata, conico-ovata, solidula, fusca, fasciis angustis albis varic ornata; spira conica, obtusa; anfr. 5 modice convexis, ultimo spiram paulo superante, basi rotundato; columella leviter arcuata; apertura parum obliqua, truncato-ovali, intus albida; perist. simplice, margine dextro recto, columellari per dilatato, umbilicum angustum fere tegente.*
 Long. 27, diam. 16 mill.
Hab. Bolivia.
20. *BULIMUS SABATIERI*, Pfr. *B. testa subperforata, ovato-oblonga, tenuiuscula, sublevigata, irregulariter striatula, strigis undulatis pallide lutescentibus et saturate brunneis alternantibus picta; spira conica, sursum pallidior, apice obtuso; anfr. 6 parum convexis, ultimo spira paulo brevior, infra medium obsolete angulato; columella stricta, violaceo-fusca; apertura parum*

obliqua, anguste elliptica; perist. simplice, recto, margine columellari sursum dilatato, reflexo, subadnato.

Long. 22, diam. 12 mill; ap. 11 mill. longa, $5\frac{1}{2}$ lata.

Hab. Banks of the "Fleuve blanc," China? (Mr. Sabatier).

21. *BULIMUS DUTAILLYI*, Pfr. *B. testa subperforata, oblongo-turrita, tenui, confertim plicatula, nitida, albida, fasciis 6-7 interruptis spadiceis ornata; spira elongata, apice acuta; anfr. 7 convexiusculis, ultimo $\frac{2}{5}$ longitudinis subæquante, basi attenuato; columella vix arcuata, subrecedente; apertura vix obliqua, elliptico-oblonga; perist. simplice, recto, margine columellari papyraceo, superne reflexo, subadnato.*

Long. 31, diam. 12 mill.; ap. $13\frac{1}{2}$ mill. longa, $6\frac{1}{2}$ lata.

Hab. Brazils (Mr. Dutailly).

22. *BULIMUS Pliculatus*, Pfr. *B. testa umbilicata, ovato-conica, tenuiuscula, plicis levibus, subvariciformibus sculpta, haud nitente, pallide grisea, strigis angustissimis albis et fuscis irregulariter picta; spira conica, acuta; anfr. 7 convexiusculis, ultimo spira paulo brevior, basi circa umbilicum medicrem, rotundum subcompresso; columella substricta; apertura vix obliqua, elliptico-oblonga; perist. simplice, recto, margine columellari a basi dilatato, subfornicatim reflexo.*

Long. 23, diam. $11\frac{1}{3}$ mill.

Hab. Bolivia.

23. *BULIMUS Clouéi*, Pfr. *B. testa perforata, ovato-turrita, solidula, striata et interdum submalleata, alba, strigis, maculis et punctis corneo-fascis irregulariter notata; spira elongato-conica, acutiuscula; anfr. 7 modice convexis, ultimo $\frac{3}{7}$ longitudinis vix æquante, subangulato, circa perforationem non perviam corneo-areolato; columella breviter recedente; apertura obliqua, sinuato-ovali, intus fuscata; perist. breviter expanso, margine columellari superne dilatato, reflexo, tum angulo obtuso ad basin descendente.*

Long. 22, diam. 10 mill.

Hab. Brazil (Mr. Cloué).

24. *BULIMUS puncticulatus*, Pfr. *B. testa profunde et compresse umbilicata, ovato-conica, tenuiuscula, sublævigata, nitida, albida, punctis pellucidis raris conspersa; spira elongato-conica, acutiuscula; anfr. 7 planiusculis, summis lutescentibus, ultimo spira brevior, antice ascendente, basi levissime compresso; columella subrecedente, leviter arcuata; apertura subverticali, oblongo-ovali; perist. simplice, margine dextro breviter expanso, columellari per dilatato, subflexuoso, patente.*

Long. 29, diam. $12\frac{1}{2}$ mill.

Hab. Bolivia.

25. *PARTULA ALABASTRINA*, Pfr. *P. testa compresse umbilicata, oblongo-conica, tenui, levissime striatula, vix nitidula, lutescenti-alabastrina; spira conica, obtusula; anfr. $5\frac{1}{2}$ convexis, ultimo*

spiram vix superante, basi attenuato, subcompresso; apertura parum obliqua, oblique truncato-oblonga; columella substrictè recedente; perist. albo, marginibus callo tenuissimo junctis, dextro late expanso, columellari latissimo, patente.

Long. 23, diam. 11 mill.

Hab. Salomon's Islands.

26. *CYCLOSTOMA (CYCLOTUS) LINDSTEDTI*, Pfr. *C. testa umbilicata, depressa, subdiscoidea, distincte striata, strigis fulgurantibus nigro-fuscis et luteis, superne latis, subtus linearibus picta; spira vix elevata; anfr. 4 convexis, celeriter accrescentibus, ultimo terete, vix descendente; umbilico lato. $\frac{1}{3}$ diametri superante; apertura obliqua, subcirculari; perist. breviter adnato, duplicato; interno expansiusculo, externo albo, undique subæqualiter patente.—Operc. Cycloti.*

Diam. maj. 11, min. 9, alt. 4 mill.

Hab. Mount Ophir, Malacca (Rev. F. W. Lindstedt).

27. *CATAULUS HÆMASTOMUS*, Pfr. *C. testa breviter et profunde rimata, ovali-pyramidali, tenuiuscula, conferte striatula, parum nitente, diaphana, citrina; spira convexiusculo-turrita, apice subacuta; sutura submarginata; anfr. 8 convexis, ultimo vix attenuato, basi axin excedente; carina umbilicali valida, compressa, antrorsum vix dilatata; periomphalo lato, costulato-striato; apertura subcirculari; perist. fusco-sanguineo, simplice, ad anfractum contiguum angustato, lateribus rectangule late patente et revoluta, basi subproducta, canali mediocri perforato.*

Long. 27-28, diam. 11-12 mill.

Hab. Ceylon (Mr. Thwaites).

28. *HELICINA (ALCADIA) RHAMPHOSTYLA*, Pfr. *H. testa conoideo-globosa, solida, sublævigata, albida vel lutescente; spira breviter conoidea, obtusula; anfr. $4\frac{1}{2}$, superis vix convexiusculis, ultimo magno, rotundato; columella lata, superne in calium crassum, semicircularem dilatata, basi dextrorsum curvata; apertura obliqua, subsemicirculari; perist. expanso, tenui, intus calloso, margine dextro subrependo, basali sinu profundo ab extremitate rostriformi columellæ disjuncto, præterea plica pone columellam intrante munito.—Operc.?*

Diam. maj. $15\frac{1}{2}$, min. 13, alt. 12 mill.

Hab. —?

29. *HELICINA NORFOLKENSIS*, Pfr. *H. testa depresso-turbinata, solidula, leviter striata, albida; spira conoidea, versus apicem minute papillarem lutescente; anfr. 5 vix convexiusculis, ultimo antice rugoso, periphæria subangulato, basi planiusculo, callo granuloso nitido circumscripto et epidermide fulva decidua obducto; columella brevi, arcuata; apertura fere diagonali, subtriangulari-lunari; perist. simplice, recto, margine basali subincrassato.—Operc. testaceum, pallidum.*

Diam. maj. 14, min. 12, alt. 8 mill.

Hab. Norfolk Islands.

30. *HELICINA PICTELLA*, Pfr. *H. testa conoideo-depressa, tenui, sub lente tenuiter et subconferte lirata, parum nitida, diaphana, pallide cornea, ad suturam rubro-maculata; spira breviter conoidea; anfr. 4 convexiusculis, ultimo subangulato, basi callo tenui subcircumscripto obducto; columella brevissima, simplice, tenui; apertura parum obliqua, semiovali; perist. simplice, recto, margine basali angulum rectum cum columella formante.—Operc.?*
 Diam. maj. 4, min. $3\frac{1}{3}$, alt. 2 mill.
Hab. Norfolk Islands.
31. *HELICINA DRAYTONENSIS*, Pfr. *H. testa conoidea, solidula, ruguloso-striata striisque spiralibus nonnullis notata, parum nitida, carnea; spira conoidea, apice mucronulata, lutea; anfr. $4\frac{1}{2}$ planiusculis, ultimo subcarinato, basi convexiore, callo tenui, subdif-fuso obducto; apertura diagonali, triangulari-semiovali; perist. albo, breviter expanso, margine basali leviter arcuato, angulatim cum columella brevi, simplice juncto.—Operc.?*
 Diam. maj. $5\frac{1}{2}$, min. $4\frac{1}{2}$, alt. $3\frac{1}{2}$ mill.
Hab. Drayton Range, North Australia.
32. *HELICINA HEATEI*, Pfr. *H. testa subdepressa, tenuiuscula, striatula et subgranulata, parum nitida, fasciis latis saturate purpureis et albis picta; spira brevissime conoidea, mucronulata; anfr. 5 vix convexiusculis, ultimo lato, periphèria subcarinato, basi circa callum fulvo-aurantiacum, circumscriptum albo; apertura perobliqua, triangulari-semiovali; columella brevi, perarcuata; perist. reflexo, late aurantiaco, margine basali levissime arcuato, immediate in callum basalem continuato.—Operc.?*
 Diam. maj. 14, min. $11\frac{1}{3}$, alt. $7\frac{1}{2}$ mill.
 β. *Lutea, spira et fascia unica anfr. ultimi supera purpureis, callo basali et peristomate igneis.*
Hab. Island of Granada, West Indies (named after R. W. Heate, Esq., Lieut.-Governor of the Island).
33. *HELICINA RUFA*, Pfr. *H. testa subconoideo-depressa, solidula, subrugoso-striata et punctulata, nitida, rufa; spira subconoideo-convexa, vertice subtili; anfr. $4\frac{1}{2}$, superis planiusculis, ultimo lato, depresso, periphèria subrotundato; apertura diagonali, late semiovali; columella verticaliter ab anfr. penultimo descendente, leviter curvata, antice in tuberculum desinente, callum basalem emittente subgranulatum, circumscriptum; perist. breviter expanso, albo-limbato.—Operc. concolor.*
 Diam. maj. 13, min. 11, alt. $6\frac{2}{3}$ mill.
 β. *Pallide straminea, perist. intus pallide aurantiaco.*
 γ. *Minor, rufa.*
 Diam. maj. 10, min. $8\frac{2}{3}$, alt. $5\frac{1}{2}$ mill.
Hab. Haiti (Mr. Sallé).



1. *Cylindrella Ghesbreght* Pf. 2. *C. turris* Pf. 3. *C. clava* Pf.
 4. 5. *Felix neta* Pf., and var. 6. 7. *H. lombi* Pf. and var.
 8. *H. Grevillei* Pf.